

RESEARCH

Open Access



# On the existence and uniqueness of a generalized solution of the Protter problem for (3 + 1)-D Keldysh-type equations

Nedyu Popivanov<sup>1\*</sup>, Tsvetan Hristov<sup>1</sup>, Aleksey Nikolov<sup>2</sup> and Manfred Schneider<sup>3</sup>

\*Correspondence: nedyu@fmi.uni-sofia.bg  
<sup>1</sup>Faculty of Mathematics and Informatics, University of Sofia, Sofia, 1164, Bulgaria  
Full list of author information is available at the end of the article

## Abstract

A (3 + 1)-dimensional boundary value problem for equations of Keldysh type (the second kind) is studied. Such problems for equations of Tricomi type (the first kind) or for the wave equation were formulated by M.H. Protter (1954) as multidimensional analogues of Darboux or Cauchy-Goursat plane problems. Now, it is well known that Protter problems are not correctly set, and they have singular generalized solutions, even for smooth right-hand sides. In this paper an analogue of the Protter problem for equations of Keldysh type is given. An appropriate generalized solution with possible singularity is defined. Results for uniqueness and existence of such a generalized solution are obtained. Some a priori estimates are stated.

**MSC:** 35D30; 35M12; 35A20

**Keywords:** weakly hyperbolic equations; boundary value problems; generalized solutions; uniqueness; behavior of solution

## 1 Introduction

In the present paper we consider an analogue of the Protter problems for (3 + 1)-D Keldysh-type equations. For  $m \in \mathbf{R}$ ,  $0 < m < 2$ , we study some boundary value problems (BVPs) for the weakly hyperbolic equation

$$L_m[u] \equiv u_{x_1x_1} + u_{x_2x_2} + u_{x_3x_3} - (t^m u_t)_t = f(x, t), \tag{1.1}$$

expressed in Cartesian coordinates  $(x, t) = (x_1, x_2, x_3, t) \in \mathbf{R}^4$  in a simply connected region

$$\Omega_m := \left\{ (x, t) : 0 < t < t_0, \frac{2}{2-m} t^{\frac{2-m}{2}} < \sqrt{x_1^2 + x_2^2 + x_3^2} < 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

bounded by the ball  $\Sigma_0 := \{(x, t) : t = 0, \sqrt{x_1^2 + x_2^2 + x_3^2} < 1\}$ , centered at the origin  $O = (0, 0, 0, 0)$  and by two characteristic surfaces of equation (1.1)

$$\Sigma_1^m := \left\{ (x, t) : 0 < t < t_0, \sqrt{x_1^2 + x_2^2 + x_3^2} = 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

$$\Sigma_2^m := \left\{ (x, t) : 0 < t < t_0, \sqrt{x_1^2 + x_2^2 + x_3^2} = \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

where  $t_0 := \left(\frac{2-m}{4}\right)^{\frac{2}{2-m}}$ .

In this work we are interested in finding sufficient conditions for the existence and uniqueness of a generalized solution of the following problem.

**Problem PK** Find a solution to equation (1.1) in  $\Omega_m$  that satisfies the boundary conditions

$$u|_{\Sigma_1^m} = 0; \quad t^m u_t \rightarrow 0, \quad \text{as } t \rightarrow +0.$$

The adjoint problem to **PK** is as follows.

**Problem PK\*** Find a solution to the self-adjoint equation (1.1) in  $\Omega_m$  that satisfies the boundary conditions

$$u|_{\Sigma_2^m} = 0; \quad t^m u_t \rightarrow 0, \quad \text{as } t \rightarrow +0.$$

First, we present a brief historical overview here and provide an extensive list of references.

Protter arrived at similar problems, but for Tricomi-type equations, while studying BVPs which describe transonic flows in fluid dynamics. It is well known that most important boundary value problems that, in the case of linear mixed-type equations, appear in hodograph plane for two-dimensional transonic potential flows are the classical Tricomi, Frankl', and Guderley-Morawetz problems. The first two for flows in nozzles and jets and the third one as an approximation in flows about airfoils. For such connections, see the paper of Morawetz [1]. About sixty years ago Murray Protter [2] stated a multidimensional variant of the famous Guderley-Morawetz plane problem for hyperbolic-elliptic equations that had been studied by Morawetz [3], Lax and Phillips [4]. This problem now is known as the Protter-Morawetz problem. A result for uniqueness was obtained by Aziz and Schneider [5] in the case of Frankl-Morawetz problem. However, the multidimensional case is rather different, and there is no general understanding of the situation. Even the question of well posedness is not completely resolved. For different statements of multidimensional Darboux-type problems or some related Protter-Morawetz problems for mixed-type equations, see [1, 6–13]. Some Tricomi problems for the Lavrent'ev-Bitsadze equation are studied in [14–16]. On the other hand, different problems for elliptic-hyperbolic equations of Keldysh type have specific applications in plasma physics, optics, and analysis on projective spaces (see Otway [17, 18] and Otway and Marini [19]). Various statements of problems for mixed equations of Tricomi or Keldysh type can be found in Oleinik and Radkevič [20], Nakhushev [21], and several applications of such problems in the study of transonic flows are described in Chen [22], Čanić and Keyfitz [23]. Let us also mention some results in the thermodynamic theory of porous elastic bodies given in [24, 25]. In order to analyze the spatial behavior of solutions, some appropriate estimates and similar procedures are used there.

In relation to the mixed-type problems, Protter also formulated and studied some BVPs in the hyperbolic part of the domain for the wave equation [26] and degenerated hyperbolic (or weakly hyperbolic) equations of Tricomi type [2]. In that case the Protter problems are multidimensional analogues of the plane Darboux or Cauchy-Goursat problems

(see Kalmenov [27] and Nakhushhev [28]). The equations are considered in  $(3 + 1)$ -D domain, bounded by two characteristic surfaces and noncharacteristic plane region. The data are prescribed on one characteristic and on a noncharacteristic boundary part. Protter considered [2, 26] Tricomi-type equations or the wave equation ( $m \in \mathbf{R}, m \geq 0$ )

$$\tilde{L}_m[u] := t^m [u_{x_1x_1} + u_{x_2x_2} + u_{x_3x_3}] - u_{tt} = f(x, t) \tag{1.2}$$

in the domain

$$\tilde{\Omega}_m := \left\{ (x_1, x_2, x_3, t) : t > 0, \frac{2}{m+2} t^{\frac{m+2}{2}} < \sqrt{x_1^2 + x_2^2 + x_3^2} < 1 - \frac{2}{m+2} t^{\frac{m+2}{2}} \right\},$$

bounded by  $\Sigma_0$  and two characteristics surfaces of (1.2)

$$\begin{aligned} \tilde{\Sigma}_1^m &= \left\{ t > 0, \sqrt{x_1^2 + x_2^2 + x_3^2} = 1 - \frac{2}{m+2} t^{\frac{m+2}{2}} \right\}, \\ \tilde{\Sigma}_2^m &= \left\{ t > 0, \sqrt{x_1^2 + x_2^2 + x_3^2} = \frac{2}{m+2} t^{\frac{m+2}{2}} \right\}. \end{aligned}$$

He proposed four problems, known now as Protter problems.

**Protter problems** Find a solution of equation (1.2) in the domain  $\tilde{\Omega}_m$  with one of the following boundary conditions:

$$\begin{aligned} P1: \quad u|_{\Sigma_0 \cup \tilde{\Sigma}_1^m} &= 0, & P1^*: \quad u|_{\Sigma_0 \cup \tilde{\Sigma}_2^m} &= 0; \\ P2: \quad u|_{\tilde{\Sigma}_1^m} &= 0, & u_t|_{\Sigma_0} &= 0, & P2^*: \quad u|_{\tilde{\Sigma}_2^m} &= 0, & u_t|_{\Sigma_0} &= 0. \end{aligned}$$

The boundary conditions in problem  $P1^*$  (respectively  $P2^*$ ) are the adjoint boundary conditions to problem  $P1$  (respectively  $P2$ ) for (1.2) in  $\tilde{\Omega}_m$ .

It turns out that instead of both boundary conditions given in problems  $P1$  on  $\tilde{\Sigma}_1^m, \Sigma_0$  and in  $P2$  on  $\tilde{\Sigma}_2^m, \Sigma_0$  for the Tricomi-type equation (1.2), in the case of Keldysh-type equation (1.1), they are reduced to only one boundary condition on the characteristic  $\Sigma_1^m$  and a condition on the growth of possible singularity of the derivative  $u_t$  as  $t \rightarrow +0$ .

We mention some known results for Protter problems in the Tricomi case that make the investigation of such problems interesting and reasonable. Garabedian [29] obtained a result for the uniqueness of classical solution to problem  $P1$  for the wave equation (i.e., equation (1.2) with  $m = 0$ ). It is interesting that contrary to their plane analogues, the 3-D Protter problems are not well posed (see [30, 31] and the monograph of Bitsadze [32]). The reason is that the adjoint homogeneous problems  $P1^*$  and  $P2^*$  have an infinite number of linearly independent nontrivial classical solutions. On the other hand, the unique generalized solutions of 3-D problems  $P1$  and  $P2$  could have strong power-type singularity on the  $\tilde{\Sigma}_2^m$  even for smooth right-hand sides (see [31, 33, 34]). Behavior of the singular solutions to 3-D problems  $P1$  and  $P2$  is studied in [35, 36]. Such results are announced for the 4-D case as well [37]. Didenko [38] studied problems  $P1$  and  $P1^*$  for the Tricomi equation ( $m = 1$ ) in the symmetric case. Aldashev [39] studied some multidimensional analogues of Protter problems for equation (1.2), but he did not mention any possible singular solutions.

These known results for Protter problems for Tricomi-type equations and many interesting applications of different boundary value problems for equations of Keldysh type motivate us to study problems  $PK$  and  $PK^*$  and to try to find out new effects that appear. In [40] ill-posedness of 3-D Protter problems for Keldysh-type equations in the frame of classical solvability is discussed, and the results for uniqueness of quasi-regular solutions are obtained. Existence and uniqueness of generalized solutions to problem  $PK$  in that case are obtained in [41], and some singular generalized solutions are announced in [42].

In [31, 33] we study Protter problems for Tricomi-type equations. For 3-D Keldysh-type equation in [43], we formulate a new Protter problem and announce some results for the existence and uniqueness of a generalized solution in the case  $0 < m < 1$ . In [44] we announce analogical results for  $(3 + 1)$ -D equations of Keldysh type in a more general case  $0 < m < 4/3$  and claim the existence of infinitely many classical smooth solutions of the  $(3 + 1)$ -D homogeneous problem  $PK^*$ . Now, in the present paper we work in the case  $0 < m < 4/3$ . Using an appropriate Riemann-Hadamard function, we find an exact integral representation of the generalized solution and prove the results announced in [44]. To avoid an infinite number of necessary conditions in the frame of classical solvability, we give a notion of a generalized solution to problem  $PK$  which can have some singularity at the point  $O$ . In order to deal successfully with the encountered difficulties for  $\varepsilon \in (0, 1)$ , we introduce the region

$$\Omega_{m,\varepsilon} := \Omega_m \cap \{|x| > \varepsilon\},$$

where  $|x| = \sqrt{x_1^2 + x_2^2 + x_3^2}$ .

We give the following definition of a generalized solution of problem  $PK$  in the case  $0 < m < 4/3$ .

**Definition 1.1** We call a function  $u(x, t)$  a generalized solution of problem  $PK$  in  $\Omega_m$ ,  $0 < m < \frac{4}{3}$ , for equation (1.1) if:

1.  $u, u_{x_j} \in C(\bar{\Omega}_m \setminus O)$ ,  $j = 1, 2, 3$ ,  $u_t \in C(\bar{\Omega}_m \setminus \bar{\Sigma}_0)$ ;
2.  $u|_{\Sigma_1^m} = 0$ ;
3. For each  $\varepsilon \in (0, 1)$  there exists a constant  $C(\varepsilon) > 0$  such that in  $\Omega_{m,\varepsilon}$

$$|u_t(x, t)| \leq C(\varepsilon)t^{-\frac{3m}{4}}; \tag{1.3}$$

4. The identity

$$\int_{\Omega_m} \{t^m u_t v_t - u_{x_1} v_{x_1} - u_{x_2} v_{x_2} - u_{x_3} v_{x_3} - f v\} dx_1 dx_2 dx_3 dt = 0 \tag{1.4}$$

holds for all  $v$  from

$$V_m := \{v(x, t) : v \in C^2(\bar{\Omega}_m), v|_{\Sigma_2^m} = 0, v \equiv 0 \text{ in a neighborhood of } O\}.$$

**Remark 1.1** We mention that all the first derivatives of the generalized solutions of 3-D Protter problems in the Tricomi case can have singularity on the boundary of the domain (see [31, 33]). Actually, this fact corresponds to the analogical situation in a 2-D case of the Darboux problem (see [27]). While in the Keldysh case, according to Definition 1.1,

the derivative  $u_t$  can be unbounded when  $t \rightarrow +0$ , but  $u_{x_1}$ ,  $u_{x_2}$  and  $u_{x_3}$  are bounded in each  $\bar{\Omega}_{m,\varepsilon}$ ,  $\varepsilon > 0$ .

In this paper, first, we prove results for the uniqueness of a generalized solution to problem  $PK$ .

**Theorem 1.1** *If  $m \in (0, \frac{4}{3})$ , then there exists at most one generalized solution of problem  $PK$  in  $\Omega_m$ .*

Further, we use the three-dimensional spherical functions  $Y_n^s(x)$  with  $n \in \mathbf{N} \cup \{0\}$ ;  $s = 1, 2, \dots, 2n + 1$ . The functions  $Y_n^s(x)$  are defined usually on the unit sphere  $S^2 := \{(x_1, x_2, x_3) : x_1^2 + x_2^2 + x_3^2 = 1\}$ , and  $Y_n^s$  form a complete orthonormal system in  $L_2(S^2)$  (see [45]). For convenience of discussions that follow, we extend the spherical functions out of  $S^2$  radially, keeping the same notation for the extended functions  $Y_n^s(x) := Y_n^s(x/|x|)$  for  $x \in \mathbf{R}^3 \setminus \{0\}$ .

Let the right-hand side function  $f(x, t)$  of equation (1.1) be fixed as a “harmonic polynomial” of order  $l$  with  $l \in \mathbf{N} \cup \{0\}$ , and it has the following representation:

$$f(x, t) = \sum_{n=0}^l \sum_{s=1}^{2n+1} f_n^s(|x|, t) Y_n^s(x) \tag{1.5}$$

with some coefficients  $f_n^s(|x|, t)$ .

In this special case we give an existence result as well.

**Theorem 1.2** *Let  $m \in (0, \frac{4}{3})$ . Suppose that the right-hand side function  $f(x, t)$  has the form (1.5) and  $f \in C^1(\bar{\Omega}_m)$ . Then the unique generalized solution  $u(x, t)$  of problem  $PK$  in  $\Omega_m$  exists and has the form*

$$u(x, t) = \sum_{n=0}^l \sum_{s=1}^{2n+1} u_n^s(|x|, t) Y_n^s(x). \tag{1.6}$$

**Remark 1.2** Actually, when the right-hand side function  $f(x, t)$  has the form (1.5) in Theorem 1.2, we find explicit representations for the functions  $u_n^s(|x|, t)$  in (1.6). These representations involve appropriate hypergeometric functions.

In the case when the right-hand side function  $f(x, t)$  has the form (1.5), we give an a priori estimate for the generalized solution of problem  $PK$  in  $\Omega_m$  as well.

**Theorem 1.3** *Let the conditions in Theorem 1.2 be fulfilled. Then the unique generalized solution of problem  $PK$  in  $\Omega_m$  has the form (1.6) and satisfies the a priori estimate*

$$|u(x, t)| \leq c \left( \max_{\bar{\Omega}_m} |f| \right) |x|^{-l-1}, \tag{1.7}$$

with a constant  $c > 0$  independent of  $f$ .

Estimate (1.7) shows the maximal order of possible singularity at point  $O$ , when the right-hand side function  $f(x, t)$  is a “harmonic polynomial” of fixed order  $l$ . We will point out

that a similar a priori estimate for generalized solutions to 3-D Protter problem  $P1$  in the Tricomi case is obtained in [36], while an estimate from below in this case is given in [31].

The present paper contains the introduction and five more sections. In Section 2, the Protter problem  $PK$  is considered in a model case when the right-hand side function  $f(x, t)$  of equation (1.1) is fixed as a “harmonic polynomial” (1.5) of order  $l$ . In that case we formulate the 2-D boundary value problems  $PK_1$  and  $PK_2$ , corresponding to the  $(3 + 1)$ -D problem  $PK$ . We give a notion for a generalized solution of Cauchy-Goursat problem  $PK_2$ , and in Section 3, using the Riemann-Hadamard function associated to this problem, we find an integral representation for a generalized solution. Further, we obtain existence and uniqueness results for a generalized solution of problem  $PK_2$ . Actually, this is the essential result in this paper and has the most difficult proof. Using the results of the previous section, in Section 4 we prove the main results in this paper, i.e., Theorem 1.1, Theorem 1.2 and Theorem 1.3. In Appendix A we give the main properties of the Riemann-Hadamard function associated to the Cauchy-Goursat problem  $PK_2$ . In Appendix B some auxiliary results, needed for the study of the generalized solution to problem  $PK_2$ , are proven.

### 2 Two-dimensional Cauchy-Goursat problems corresponding to problem $PK$

Using spherical coordinates  $(r, \theta, \varphi, t) \in \mathbf{R}^4, 0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, r > 0$  with

$$x_1 = r \sin \theta \cos \varphi, \quad x_2 = r \sin \theta \sin \varphi, \quad x_3 = r \cos \theta$$

problem  $PK$  can suitably be treated. Written in the new coordinates, equation (1.1) becomes

$$L_m u = \frac{1}{r^2} (r^2 u_r)_r + \frac{1}{r^2 \sin \theta} (\sin \theta u_\theta)_\theta + \frac{1}{r^2 \sin^2 \theta} u_{\varphi\varphi} - (t^m u_t)_t = f. \tag{2.1}$$

We consider equation (2.1) in the region

$$\Omega_m = \left\{ (r, \theta, \varphi, t) : 0 < t < t_0, 0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, \frac{2}{2-m} t^{\frac{2-m}{2}} < r < 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

bounded by the following surfaces:

$$\begin{aligned} \Sigma_0 &= \{ (r, \theta, \varphi, t) : t = 0, 0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, r < 1 \}, \\ \Sigma_1^m &= \left\{ (r, \theta, \varphi, t) : t > 0, 0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, r = 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\}, \\ \Sigma_2^m &= \left\{ (r, \theta, \varphi, t) : t > 0, 0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, r = \frac{2}{2-m} t^{\frac{2-m}{2}} \right\}. \end{aligned}$$

Problem  $PK$  becomes the following one: find a solution to equation (2.1) with the boundary conditions

$$u|_{\Sigma_1^m} = 0; \quad t^m u_t \rightarrow 0, \quad \text{as } t \rightarrow +0.$$

The two-dimensional spherical functions, expressed in terms of  $\theta$  and  $\varphi$  in the traditional definition (see [45]), are  $Y_n^s(\theta, \varphi) := Y_n^s(x), x \in S^2, n \in \mathbf{N} \cup \{0\}, s = 1, 2, \dots, 2n + 1$ , and satisfy

the differential equation

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} Y_n^s \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} Y_n^s + n(n+1) Y_n^s = 0.$$

In the special case when the right-hand side function  $f(x, t)$  of equation (2.1) has the form

$$f(r, \theta, \varphi, t) = f_n^s(r, t) Y_n^s(\theta, \varphi),$$

we may look for a solution of the form

$$u(r, \theta, \varphi, t) = u_n^s(r, t) Y_n^s(\theta, \varphi)$$

with unknown coefficient  $u_n^s(r, t)$ .

For the coefficients  $u_n^s(r, t)$  which correspond to the right-hand sides  $f_n^s(r, t)$ , we obtain the 2-D equation

$$u_{rr} + \frac{2}{r} u_r - (t^m u_t)_t - \frac{n(n+1)}{r^2} u = f$$

in the domain

$$G_m = \left\{ (r, t) : 0 < t < t_0, \frac{2}{2-m} t^{\frac{2-m}{2}} < r < 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

which is bounded by the segment  $S_0 = \{(r, t) : 0 < r < 1, t = 0\}$  and the characteristics

$$S_1^m := \left\{ (r, t) : 0 < t < t_0, r = 1 - \frac{2}{2-m} t^{\frac{2-m}{2}} \right\},$$

$$S_2^m := \left\{ (r, t) : 0 < t < t_0, r = \frac{2}{2-m} t^{\frac{2-m}{2}} \right\}.$$

In this case, for  $u(r, t)$ , the 2-D problem corresponding to  $PK$  is the problem

$$PK_1 \quad \begin{cases} u_{rr} + \frac{2}{r} u_r - (t^m u_t)_t - \frac{n(n+1)}{r^2} u = f(r, t) & \text{in } G_m, \\ u|_{S_1^m} = 0; \quad t^m u_t \rightarrow 0, & \text{as } t \rightarrow +0. \end{cases}$$

**Remark 2.1** When the right-hand side function  $f(x, t)$  has the form (1.5), it is enough to take test functions  $v \in V_m$  in the identity (1.4) to have the form  $v = w(|x|, t) Y_n^s(x)$ ,  $n \in \mathbf{N} \cup \{0\}$ ,  $s = 1, 2, \dots, 2n + 1$  and

$$w \in V_m^{(1)} := \{w(r, t) : w \in C^2(\bar{G}_m), w|_{S_2^m} = 0, w \equiv 0 \text{ in a neighborhood of } (0, 0)\}.$$

The generalized solution of problem  $PK_1$  is defined as follows.

**Definition 2.1** We call a function  $u(r, t)$  a generalized solution of problem  $PK_1$  in  $G_m$  ( $0 < m < \frac{4}{3}$ ) if:

1.  $u, u_r \in C(\bar{G}_m \setminus (0, 0)), u_t \in C(\bar{G}_m \setminus \bar{S}_0);$
2.  $u|_{S_1^m} = 0;$
3. For each  $\varepsilon \in (0, 1)$  there exists a constant  $C(\varepsilon) > 0$  such that the estimates

$$|u_t(r, t)| \leq C(\varepsilon)t^{-\frac{3m}{4}}$$

hold in  $G_{m,\varepsilon} := G_m \cap \{r > \varepsilon\};$

4. The identity

$$\int_{G_m} \left\{ u_r v_r - t^m u_t v_t + \frac{n(n+1)}{r^2} uv + fv \right\} r^2 dr dt = 0 \tag{2.2}$$

holds for all  $v \in V_m^{(1)}.$

Substituting the new characteristic coordinates

$$\xi = 1 - r - \frac{2}{2-m}t^{\frac{2-m}{2}}, \quad \eta = 1 - r + \frac{2}{2-m}t^{\frac{2-m}{2}} \tag{2.3}$$

and the new functions

$$\begin{aligned} U(\xi, \eta) &= r(\xi, \eta)u(r(\xi, \eta), t(\xi, \eta)), \\ V(\xi, \eta) &= r(\xi, \eta)v(r(\xi, \eta), t(\xi, \eta)), \\ F(\xi, \eta) &= \frac{1}{8}(2 - \xi - \eta)f(r(\xi, \eta), t(\xi, \eta)), \end{aligned} \tag{2.4}$$

from problem  $PK_1,$  we get the 2-D Cauchy-Goursat problem

$$PK_2 : \begin{cases} U_{\xi\eta} + \frac{\beta}{\eta-\xi}(U_\xi - U_\eta) - \frac{n(n+1)}{(2-\xi-\eta)^2}U = F(\xi, \eta) & \text{in } D, \\ U(0, \eta) = 0, \quad \lim_{\eta-\xi \rightarrow +0} (\eta - \xi)^{2\beta}(U_\xi - U_\eta) = 0, \end{cases} \tag{2.5}$$

where

$$D := \{(\xi, \eta) : 0 < \xi < \eta < 1\} \subset \mathbf{R}^2,$$

and the parameter  $\beta = \frac{m}{2(2-m)} \in (0, 1)$  since  $0 < m < \frac{4}{3}.$

The generalized solution of problem  $PK_2$  is defined as follows.

**Definition 2.2** We call a function  $U(\xi, \eta)$  a generalized solution of problem  $PK_2$  in  $D$  ( $0 < \beta < 1$ ) if:

1.  $U, U_\xi + U_\eta \in C(\bar{D} \setminus (1, 1)), U_\xi - U_\eta \in C(\bar{D} \setminus \{\eta = \xi\});$
2.  $U(0, \eta) = 0;$  (2.7)

3. For each  $\varepsilon \in (0, 1)$  there exists a constant  $C(\varepsilon) > 0$  such that

$$|(U_\xi - U_\eta)(\xi, \eta)| \leq C(\varepsilon)(\eta - \xi)^{-\beta} \quad \text{in } \bar{D}_\varepsilon \setminus \{\eta = \xi\}, \tag{2.8}$$

where  $D_\varepsilon := D \cap \{\xi < 1 - \varepsilon\};$



4. The identity

$$\int_D (\eta - \xi)^{2\beta} \left\{ U_\xi V_\eta + U_\eta V_\xi + \frac{2n(n+1)}{(2-\xi-\eta)^2} UV + 2FV \right\} d\xi d\eta = 0 \tag{2.9}$$

holds for all

$$V \in V^{(2)} := \{ V(\xi, \eta) : V \in C^2(\bar{D}), V(\xi, 1) = 0, V \equiv 0 \text{ in a neighborhood of } (1, 1) \}.$$

### 3 Existence and uniqueness of a generalized solution to the Cauchy-Goursat plane problem $PK_2$

In this section we prove the existence and uniqueness of a generalized solution to problem  $PK_2$ . In order to do this, we use the Riemann-Hadamard function associated to problem  $PK_2$  to find an integral representation for a generalized solution of this problem in  $D$ . According to Gellerstedt [46] and the results of Nakhushev mentioned in the book of Smirnov [47], this function has the form

$$\Phi(\xi, \eta; \xi_0, \eta_0) = \begin{cases} \Phi^+(\xi, \eta; \xi_0, \eta_0), & \eta > \xi_0, \\ \Phi^-(\xi, \eta; \xi_0, \eta_0), & \eta < \xi_0, \end{cases} \tag{3.1}$$

for  $(\xi_0, \eta_0) \in D$  and  $(\xi, \eta) \in \bar{T} \cup \bar{\Pi} \setminus \{\eta = \xi_0\}$ , where

$$T := \{(\xi, \eta) : 0 < \xi < \eta < \xi_0\}, \quad \Pi := \{(\xi, \eta) : 0 < \xi < \xi_0, \xi_0 < \eta < \eta_0\}.$$

The Riemann-Hadamard function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  should have the following main properties (see [46, 47]):

(i) The function  $\Phi$  as a function of  $(\xi_0, \eta_0)$  satisfies

$$\begin{aligned} E_{\xi_0, \eta_0}[\Phi] &:= \frac{\partial^2 \Phi}{\partial \xi_0 \partial \eta_0} + \frac{\beta}{\eta_0 - \xi_0} \left( \frac{\partial \Phi}{\partial \xi_0} - \frac{\partial \Phi}{\partial \eta_0} \right) - \frac{n(n+1)}{(2-\xi_0-\eta_0)^2} \Phi \\ &= 0 \quad \text{in } D, \eta \neq \xi_0 \end{aligned} \tag{3.2}$$

and with respect to the first pair of variables  $(\xi, \eta)$

$$\begin{aligned} E_{\xi, \eta}^*[\Phi] &:= \frac{\partial^2 \Phi}{\partial \xi \partial \eta} - \frac{\partial}{\partial \xi} \left( \frac{\beta \Phi}{\eta - \xi} \right) + \frac{\partial}{\partial \eta} \left( \frac{\beta \Phi}{\eta - \xi} \right) - \frac{n(n+1)}{(2-\xi-\eta)^2} \Phi \\ &= 0 \quad \text{in } D, \eta \neq \xi_0; \end{aligned} \tag{3.3}$$

(ii)  $\Phi^+(\xi_0, \eta_0; \xi_0, \eta_0) = 1$ ;

(iii)  $\Phi^+(\xi, \eta_0; \xi_0, \eta_0) = \left(\frac{\eta_0 - \xi}{\eta_0 - \xi_0}\right)^\beta$ ;

(iv)  $\Phi^+(\xi_0, \eta; \xi_0, \eta_0) = \left(\frac{\eta - \xi_0}{\eta_0 - \xi_0}\right)^\beta$ ;

(v) The jump of the function  $\Phi$  on the line  $\{\eta = \xi_0\}$  is

$$\begin{aligned} [[\Phi]] &:= \lim_{\delta \rightarrow +0} \{ \Phi^-(\xi, \xi_0 - \delta; \xi_0, \eta_0) - \Phi^+(\xi, \xi_0 + \delta; \xi_0, \eta_0) \} \\ &= \cos(\pi\beta) \lim_{\delta \rightarrow +0} \{ \Phi^+(\xi, \xi_0 + \delta; \xi_0, \xi_0 + \delta) \Phi^+(\xi_0, \xi_0 + \delta; \xi_0, \eta_0) \} \\ &= \cos(\pi\beta) \left( \frac{\xi_0 - \xi}{\eta_0 - \xi_0} \right)^\beta; \end{aligned}$$

(vi)  $\Phi^-$  vanishes on the line  $\{\eta = \xi\}$  of power  $2\beta$ .

Actually, the function  $\Phi^+$  is the Riemann function for equation (2.5).

**Remark 3.1** In the case  $0 < \beta < 1/2$  and  $F(\xi, \eta) = (\eta - \xi)^{-4\beta} f(\xi, \eta)$ , where  $f \in C(\bar{D})$ , a generalized solution of problem  $PK_2$  has an explicit integral representation (see [46] and [47]). We find an integral representation in the case  $0 < \beta < 1$  and  $F \in C(\bar{D})$  using the properties of the Riemann-Hadamard function  $\Phi(\xi, \eta; \xi_0, \eta_0)$ . The existence of a function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  with properties (i) ÷ (vi) is shown in Appendix A (see also [44]).

**Theorem 3.1** *Let  $0 < \beta < 1$  and  $F \in C(\bar{D})$ . Then each generalized solution of problem  $PK_2$  in  $D$  has the following integral representation:*

$$U(\xi_0, \eta_0) = \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) \Phi(\xi, \eta; \xi_0, \eta_0) d\eta d\xi. \tag{3.4}$$

*Proof* Let  $U(\xi, \eta)$  be a generalized solution of problem  $PK_2$  in  $D$ . For any arbitrary function  $\psi(\xi, \eta)$  from  $C_0^\infty(D)$ , we have  $\psi \in V^{(2)}$ , and from (2.9) we obtain the identity

$$\int_D (\eta - \xi)^{2\beta} \left\{ U_{\xi\eta} + \frac{\beta}{\eta - \xi} (U_{\xi\xi} - U_{\eta\eta}) - \frac{n(n+1)}{(2 - \xi - \eta)^2} U - F \right\} \psi d\xi d\eta = 0,$$

where  $U_{\xi\eta}$  is the weak derivative of  $U$ . Therefore

$$U_{\xi\eta} = F + \frac{n(n+1)}{(2 - \xi - \eta)^2} U - \frac{\beta}{\eta - \xi} (U_{\xi\xi} - U_{\eta\eta}) \in C(D)$$

since  $F, U, U_{\xi\xi} - U_{\eta\eta} \in C(D)$ . From this it follows that  $U_{\xi\eta}$  is a classical derivative of  $U$  and  $U(\xi, \eta)$  satisfies the differential equation (2.5) in  $D$  in a classical sense.

Now, using the properties of the Riemann-Hadamard function  $\Phi(\xi, \eta; \xi_0, \eta_0)$ , we obtain the integral representation (3.4) for a generalized solution of problem  $PK_2$  by integrating the identity

$$\Phi(\xi, \eta; \xi_0, \eta_0) E_{\xi, \eta} [U(\xi, \eta)] - U(\xi, \eta) E_{\xi, \eta}^* [\Phi(\xi, \eta; \xi_0, \eta_0)] = F(\xi, \eta) \Phi(\xi, \eta; \xi_0, \eta_0)$$

over a triangle

$$T_\delta := \{(\xi, \eta) : 0 < \xi < \xi_0 - 2\delta, \xi + \delta < \eta < \xi_0 - \delta\}$$

and then over the rectangle

$$\Pi_\delta := \{(\xi, \eta) : 0 < \xi < \xi_0 - 2\delta, \xi_0 + \delta < \eta < \eta_0\}$$

with  $\delta > 0$  small enough, and finally letting  $\delta \rightarrow 0$ . □

Theorem 3.1 claims the uniqueness of a generalized solution to problem  $PK_2$ . Next, we prove that if  $F \in C^1(\bar{D})$  and  $U(\xi, \eta)$  is a function defined by (3.4) in  $D$ , then  $U(\xi, \eta)$  is a generalized solution to problem  $PK_2$  in  $D$ . In order to do this, we introduce the notation

$$M_F := \max \left\{ \max_{\bar{D}_0} |F|, \max_{\bar{D}_0} |F_\xi + F_\eta| \right\},$$

and we mention that, according to Lemma A.1 (see Appendix A below), the Riemann-Hadamard function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  can be decomposed in the following way:

$$\Phi(\xi, \eta; \xi_0, \eta_0) = H(\xi, \eta; \xi_0, \eta_0) + G(\xi, \eta; \xi_0, \eta_0),$$

where  $H(\xi, \eta; \xi_0, \eta_0)$  is the Riemann-Hadamard function (A.12) associated to problem  $PK_2$  in the case  $n = 0$  and  $G(\xi, \eta; \xi_0, \eta_0)$  is an additional term. Therefore we can rewrite representation (3.4) in the form

$$U(\xi_0, \eta_0) = U^H(\xi_0, \eta_0) + U^G(\xi_0, \eta_0), \tag{3.5}$$

where

$$U^H(\xi_0, \eta_0) := \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) H(\xi, \eta; \xi_0, \eta_0) d\eta d\xi \tag{3.6}$$

and

$$U^G(\xi_0, \eta_0) := \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) G(\xi, \eta; \xi_0, \eta_0) d\eta d\xi. \tag{3.7}$$

Firstly, we will study the function  $U^H(\xi_0, \eta_0)$ . To do this, we use the estimates for some integrals involving function  $H(\xi, \eta; \xi_0, \eta_0)$  obtained in Appendix B.

**Theorem 3.2** *Let  $0 < \beta < 1$  and  $F \in C^1(\bar{D})$ . Then, for the function  $U^H(\xi_0, \eta_0)$ , we have  $U^H, U_{\xi_0}^H + U_{\eta_0}^H \in C(\bar{D} \setminus (1, 1))$ ,  $U_{\eta_0}^H \in C(\bar{D} \setminus \{\eta_0 = \xi_0\})$  and the following estimates hold:*

$$\begin{aligned} |U^H(\xi_0, \eta_0)| &\leq K_1 M_F \xi_0 \quad \text{in } \bar{D} \setminus (1, 1), \\ |U_{\xi_0}^H + U_{\eta_0}^H|(\xi_0, \eta_0) &\leq K_1 M_F \eta_0 \quad \text{in } \bar{D} \setminus (1, 1), \\ |U_{\eta_0}^H(\xi_0, \eta_0)| &\leq K_1 M_F \xi_0 (\eta_0 - \xi_0)^{-\beta} \quad \text{in } \bar{D} \setminus \{\eta_0 = \xi_0\}, \end{aligned}$$

where  $K_1 > 0$  is a constant independent of  $F$ .

*Proof Step 1.* From (3.6) and (B.1) from Lemma B.1 (see Appendix B) we obtain

$$|U^H(\xi_0, \eta_0)| \leq M_F \int_0^{\xi_0} \int_{\xi}^{\eta_0} H(\xi, \eta; \xi_0, \eta_0) d\eta d\xi = M_F I^1(\xi_0, \eta_0) \leq k_1 M_F \xi_0.$$

*Step 2.* Differentiating (3.6) with respect to  $\eta_0$  and using (B.4) from Lemma B.2, we obtain

$$\begin{aligned} |U_{\eta_0}^H(\xi_0, \eta_0)| &= \left| \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) H_{\eta_0}(\xi, \eta; \xi_0, \eta_0) d\eta d\xi + \int_0^{\xi_0} F(\xi, \eta_0) \frac{(\eta_0 - \xi)^\beta}{(\eta_0 - \xi_0)^\beta} d\xi \right| \\ &\leq M_F \left\{ \int_0^{\xi_0} \int_{\xi}^{\eta_0} |H_{\eta_0}(\xi, \eta; \xi_0, \eta_0)| d\eta d\xi + \int_0^{\xi_0} \left( \frac{\eta_0 - \xi}{\eta_0 - \xi_0} \right)^\beta d\xi \right\} \\ &= M_F \left\{ I^2(\xi_0, \eta_0) + \int_0^{\xi_0} \left( \frac{\eta_0 - \xi}{\eta_0 - \xi_0} \right)^\beta d\xi \right\} \\ &\leq M_F (k_2 + 1) \xi_0 (\eta_0 - \xi_0)^{-\beta}. \end{aligned}$$

*Step 3.* According to Remark A.1, the derivatives  $H_{\xi_0}^+$ ,  $H_{\xi_0}^-$  have singularities of order  $|\eta - \xi_0|^{-1}$  on the line  $\{\eta = \xi_0\}$ . Gellerstedt [46] and Moiseev [48] consider the case  $n = 0$  and suggest differentiating (3.6) after appropriate substitutions of variables. In that way one can find integral representations for the first derivatives of the solution which do not involve the first derivatives of function  $H$ . In order to do this, following Moiseev [48], we introduce new variables

$$\tilde{\xi} := \frac{\xi_0 - \xi}{\eta_0 - \xi_0}, \quad \tilde{\eta} := \frac{\eta_0 - \eta}{\eta_0 - \xi_0}. \tag{3.8}$$

We define

$$\tilde{H}^+(\tilde{\xi}, \tilde{\eta}) := H^+(\xi, \eta; \xi_0, \eta_0), \quad \tilde{H}^-(\tilde{\xi}, \tilde{\eta}) := H^-(\xi, \eta; \xi_0, \eta_0),$$

from (A.12) we obtain

$$\begin{aligned} \tilde{H}^+(\tilde{\xi}, \tilde{\eta}) &= (1 - \tilde{\eta} + \tilde{\xi})^\beta F\left(\beta, 1 - \beta, 1; \frac{\tilde{\xi}\tilde{\eta}}{1 - \tilde{\eta} + \tilde{\xi}}\right), \quad \tilde{\eta} < 1, \\ \tilde{H}^-(\tilde{\xi}, \tilde{\eta}) &= \frac{k(1 - \tilde{\eta} + \tilde{\xi})^{2\beta}}{\tilde{\xi}^\beta \tilde{\eta}^\beta} F\left(\beta, \beta, 2\beta; \frac{1 - \tilde{\eta} + \tilde{\xi}}{\tilde{\xi}\tilde{\eta}}\right), \quad \tilde{\eta} > 1. \end{aligned}$$

Then we have

$$\begin{aligned} U^H(\xi_0, \eta_0) &= (\eta_0 - \xi_0)^2 \int_0^{\frac{\xi_0}{\eta_0 - \xi_0}} \left\{ \int_0^{1 + \tilde{\xi}} F(\xi_0 - (\eta_0 - \xi_0)\tilde{\xi}, \eta_0 - (\eta_0 - \xi_0)\tilde{\eta}) \tilde{H}(\tilde{\xi}, \tilde{\eta}) d\tilde{\eta} \right\} d\tilde{\xi}, \end{aligned}$$

and

$$\begin{aligned} (U_{\xi_0}^H + U_{\eta_0}^H)(\xi_0, \eta_0) &= (\eta_0 - \xi_0)^2 \int_0^{\frac{\xi_0}{\eta_0 - \xi_0}} \int_0^{1 + \tilde{\xi}} (F_\xi + F_\eta)(\xi_0 - (\eta_0 - \xi_0)\tilde{\xi}, \eta_0 - (\eta_0 - \xi_0)\tilde{\eta}) \tilde{H}(\tilde{\xi}, \tilde{\eta}) d\tilde{\eta} d\tilde{\xi} \\ &\quad + (\eta_0 - \xi_0) \int_0^{\frac{\eta_0}{\eta_0 - \xi_0}} F(0, \eta_0 - (\eta_0 - \xi_0)\tilde{\eta}) \tilde{H}\left(\frac{\xi_0}{\eta_0 - \xi_0}, \tilde{\eta}\right) d\tilde{\eta}. \end{aligned}$$

Now the inverse transform of (3.8) gives

$$\begin{aligned} (U_{\xi_0}^H + U_{\eta_0}^H)(\xi_0, \eta_0) &= \int_0^{\xi_0} \int_\xi^{\eta_0} (F_\xi + F_\eta)(\xi, \eta) H(\xi, \eta; \xi_0, \eta_0) d\eta d\xi \\ &\quad + \int_0^{\eta_0} F(0, \eta) H(0, \eta; \xi_0, \eta_0) d\eta. \end{aligned}$$

Now (B.1) from Lemma B.1 and (B.6) from Lemma B.3 give

$$|(U_{\xi_0}^H + U_{\eta_0}^H)(\xi_0, \eta_0)| \leq M_F \{I^1(\xi_0, \eta_0) + I^3(\xi_0, \eta_0)\} \leq M_F(k_1 + k_3)\eta_0. \quad \square$$

**Theorem 3.3** *Let the conditions in Theorem 3.2 be fulfilled. Then for the function  $U^G(\xi_0, \eta_0)$  we have  $U^G, U_{\xi_0}^G, U_{\eta_0}^G \in C(\bar{D} \setminus (1, 1))$ , and the following estimates hold in  $\bar{D} \setminus (1, 1)$ :*

$$|U^G(\xi_0, \eta_0)| \leq K_2 M_F \xi_0 (2 - \xi_0 - \eta_0)^{-n}, \tag{3.9}$$

$$|U_{\xi_0}^G(\xi_0, \eta_0)| \leq K_2 M_F \xi_0 (2 - \xi_0 - \eta_0)^{-n-1}, \tag{3.10}$$

$$|U_{\eta_0}^G(\xi_0, \eta_0)| \leq K_2 M_F \xi_0 (2 - \xi_0 - \eta_0)^{-n-1}, \tag{3.11}$$

where  $K_2 > 0$  is a constant independent of  $F$ .

*Proof* Using estimates (A.26) and (A.27), from (3.7) we obtain estimate (3.9):

$$\begin{aligned} |U^G(\xi_0, \eta_0)| &= \left| \int_0^{\xi_0} \int_{\xi}^{\xi_0} F(\xi, \eta) G^-(\xi, \eta; \xi_0, \eta_0) d\eta d\xi \right. \\ &\quad \left. + \int_0^{\xi_0} \int_{\xi_0}^{\eta_0} F(\xi, \eta) G^+(\xi, \eta; \xi_0, \eta_0) d\eta d\xi \right| \\ &\leq C_G M_F \xi_0 \left\{ \frac{1}{2} \xi_0 (2 - \xi_0 - \eta_0)^{-n} + (\eta_0 - \xi_0)^{1-\beta} \right\} \\ &\leq K_2 M_F \xi_0 (2 - \xi_0 - \eta_0)^{-n}. \end{aligned}$$

Now we calculate

$$U_{\xi_0}^G(\xi_0, \eta_0) = \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) G_{\xi_0}(\xi, \eta; \xi_0, \eta_0) d\eta d\xi.$$

Here we do not have integrals on the boundaries because  $Y = 0$  on the line  $\{\xi = \xi_0\}$ , and the function  $G(\xi, \eta, \xi_0, \eta_0)$  has no jump on the line  $\{\eta = \xi_0\}$  (see Appendix A). Applying estimates (A.30) and (A.31) to this integral, we have

$$\begin{aligned} |U_{\xi_0}^G(\xi_0, \eta_0)| &\leq \frac{M_F C_G}{(2 - \xi_0 - \eta_0)^{n+1}} \int_0^{\xi_0} \int_{\xi}^{\xi_0} (\xi_0 - \eta)^{-\beta} d\eta d\xi \\ &\quad + \frac{M_F C_G}{2 - \xi_0 - \eta_0} \int_0^{\xi_0} \int_{\xi_0}^{\eta_0} (\eta - \xi_0)^{-\beta} d\eta d\xi \\ &\leq \frac{M_F C_G}{(2 - \xi_0 - \eta_0)^{n+1}} (I_1^1 + 2^n I_2^1). \end{aligned}$$

Now (B.2) and (B.3) from Lemma B.1 give estimate (3.10). Further, we calculate

$$U_{\eta_0}^G(\xi_0, \eta_0) = \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) G_{\eta_0}(\xi, \eta; \xi_0, \eta_0) d\eta d\xi,$$

where we used that  $Y = 0$  on the line  $\eta = \eta_0$ . Analogously, applying estimates (A.28) and (A.29), which are even better than (A.30) and (A.31), to the last integral for the derivative  $G_{\eta_0}$ , we obtain estimate (3.11). □

As a direct consequence of Theorem 3.2 and Theorem 3.3, in view of  $U = U^H + U^G$ , we have the following theorem.

**Theorem 3.4** *Let  $0 < \beta < 1$  and  $F \in C^1(\bar{D})$ . Then, for the function  $U(\xi, \eta)$  defined by (3.4), we have  $U, U_\xi + U_\eta \in C(\bar{D} \setminus (1, 1))$ ,  $U_\eta \in C(\bar{D} \setminus \{\eta = \xi\})$  and for some constant  $K_3 > 0$  the estimates below hold*

$$\begin{aligned} |U(\xi, \eta)| &\leq K_3 M_F \xi (2 - \xi - \eta)^{-n} \quad \text{in } \bar{D} \setminus (1, 1), \\ |(U_\xi + U_\eta)(\xi, \eta)| &\leq K_3 M_F (2 - \xi - \eta)^{-n-1} \quad \text{in } \bar{D} \setminus (1, 1), \\ |U_\eta(\xi, \eta)| &\leq K_3 M_F \xi (\eta - \xi)^{-\beta} (2 - \xi - \eta)^{-n-1} \quad \text{in } \bar{D} \setminus \{\eta = \xi\}. \end{aligned} \tag{3.12}$$

Now, we are able to prove the following existence result.

**Theorem 3.5** *Let  $0 < \beta < 1$  and  $F \in C^1(\bar{D})$ . Then there exists one and only one generalized solution to problem  $PK_2$  in  $D$ , which has integral representation (3.4), and it satisfies estimates (3.12).*

*Proof* Let  $U(\xi, \eta)$  be the function known from Theorem 3.4. Therefore  $U, U_\xi + U_\eta \in C(\bar{D} \setminus (1, 1))$ ,  $U_\eta \in C(\bar{D} \setminus \{\eta = \xi\})$ , and it satisfies estimates (3.12) in Definition 2.2. But in view of (3.12) it is obvious that condition (2.7) and estimate (2.8) hold.

To prove that  $U(\xi, \eta)$  satisfies identity (2.9) in Definition 2.2, we need several steps as follows.

*Step 1.* We prove that  $U(\xi, \eta)$  satisfies the differential equation (2.5) in a classical sense and  $\frac{\partial}{\partial \eta}(U_\xi) \in C(D)$ .

(1.i) Following Smirnov [47], we find another representation formula for the function  $U^H(\xi, \eta)$ . Let us introduce the function

$$R_0(\xi, \eta; \xi_0, \eta_0) := \begin{cases} R_0^+(\xi, \eta; \xi_0, \eta_0), & \eta > \xi_0, \\ R_0^-(\xi, \eta; \xi_0, \eta_0), & \eta < \xi_0, \end{cases}$$

where

$$\begin{aligned} R_0^+(\xi, \eta; \xi_0, \eta_0) &:= \left(\frac{\eta_0 - \eta}{\eta_0 - \xi_0}\right)^\beta \left(\frac{\eta_0 - \eta}{\eta_0 - \xi}\right)^{1-\beta} F_1\left(1 - \beta, \beta, 1 - \beta, 2; \frac{\eta_0 - \eta}{\eta_0 - \xi_0}, \frac{\eta_0 - \eta}{\eta_0 - \xi}\right), \\ R_0^-(\xi, \eta; \xi_0, \eta_0) &:= \gamma \left(\frac{\eta - \xi}{\xi_0 - \xi}\right)^\beta \left(\frac{\eta - \xi}{\eta_0 - \xi}\right)^\beta F_1\left(\beta, \beta, \beta, 1 + 2\beta; \frac{\eta - \xi}{\xi_0 - \xi}, \frac{\eta - \xi}{\eta_0 - \xi}\right). \end{aligned}$$

Here  $\gamma = -\frac{\Gamma(\beta)}{\Gamma(1-\beta)\Gamma(1+2\beta)}$  and  $F_1(a, b_1, b_2, c; x, y)$  is the hypergeometric function (A.8) of two variables (see Appendix A).

In [47] the case  $0 < \beta < 1/2$  is considered, but here we find that in a more general case  $0 < \beta < 1$  the function  $R_0(\xi, \eta; \xi_0, \eta_0)$  solves

$$\begin{aligned} \frac{\partial R_0}{\partial \eta} &= -(\eta - \xi)^{-1} H(\xi, \eta; \xi_0, \eta_0) \quad \text{for } (\xi, \eta) \in \Pi \cup T, \\ R_0|_{\eta=\eta_0} &= 0, \quad R_0|_{\eta=\xi} = 0, \end{aligned} \tag{3.13}$$

where  $(\xi_0, \eta_0) \in D$  and  $H(\xi, \eta; \xi_0, \eta_0)$  is function (A.12).

Using (3.13), integration by parts and

$$[R_0^+ - R_0^-]|_{\eta=\xi_0} = \frac{1}{\beta}$$

leads to the integral representation

$$\begin{aligned}
 U^H(\xi_0, \eta_0) &:= \int_0^{\xi_0} \int_{\xi}^{\eta_0} \frac{\partial}{\partial \eta} [(\eta - \xi)F(\xi, \eta)] R_0(\xi, \eta; \xi_0, \eta_0) \, d\eta \, d\xi \\
 &\quad + \frac{1}{\beta} \int_0^{\xi_0} (\xi_0 - \xi)F(\xi, \xi_0) \, d\xi.
 \end{aligned}
 \tag{3.14}$$

(1.ii) Differentiating (3.14) we obtain that  $U^H$  satisfies the differential equation

$$(U^H_{\xi_0})_{\eta_0} + \frac{\beta}{\eta_0 - \xi_0} (U^H_{\xi_0} - U^H_{\eta_0}) = F(\xi_0, \eta_0),
 \tag{3.15}$$

where all derivatives are in a classical sense and they are continuous in  $D$ .

(1.iii) Since  $H(\xi, \eta; \xi_0, \eta_0)$  satisfies the differential equation (3.2) with  $n = 0$  and  $\Phi = H + G$  satisfies (3.2) with  $n \geq 0$  for the difference  $G = \Phi - H$ , we obtain

$$G_{\xi_0 \eta_0} + \frac{\beta}{\eta_0 - \xi_0} (G_{\xi_0} - G_{\eta_0}) - \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} G = \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} H.$$

Now, using integral representation (3.7) for  $U^G(\xi_0, \eta_0)$ , we calculate

$$\begin{aligned}
 (U^G_{\xi_0})_{\eta_0} &+ \frac{\beta}{\eta_0 - \xi_0} (U^G_{\xi_0} - U^G_{\eta_0}) - \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} U^G \\
 &= \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) \left[ G_{\xi_0 \eta_0} + \frac{\beta}{\eta_0 - \xi_0} (G_{\xi_0} - G_{\eta_0}) \right. \\
 &\quad \left. - \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} G \right] (\xi, \eta; \xi_0, \eta_0) \, d\eta \, d\xi \\
 &= \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} \int_0^{\xi_0} \int_{\xi}^{\eta_0} F(\xi, \eta) H(\xi, \eta; \xi_0, \eta_0) \, d\eta \, d\xi \\
 &= \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} U^H,
 \end{aligned}
 \tag{3.16}$$

where all derivatives are in a classical sense and they are continuous in  $D$ .

(1.iv) Since  $U = U^H + U^G$ , summing up equations (3.15) and (3.16), we obtain the differential equation

$$(U_{\xi_0})_{\eta_0} + \frac{\beta}{\eta_0 - \xi_0} (U_{\xi_0} - U_{\eta_0}) - \frac{n(n+1)}{(2 - \xi_0 - \eta_0)^2} U = F(\xi_0, \eta_0)$$

in a classical sense. But, since  $F, U, U_{\xi_0} - U_{\eta_0} \in C(D)$ , it follows that  $(U_{\xi_0})_{\eta_0} \in C(D)$ .

Step 2. We will prove that identity (2.9) holds for all  $V(\xi, \eta) \in V^{(2)}$ .

(2.i) Let  $V(\xi, \eta) \in V^{(2)}$  and in addition  $V(\xi, \eta) \equiv 0$  in a neighborhood of  $\{\eta = \xi\}$  and in a neighborhood of  $\{\eta = 1\}$ . From Step 1 we know that  $U(\xi, \eta)$  satisfies the differential equation (2.5), where all derivatives are in a classical sense, continuous in  $D$ . Let us consider

$$I_V := \int_D (\eta - \xi)^{2\beta} \left\{ U_{\xi} V_{\eta} + U_{\eta} V_{\xi} + \frac{2n(n+1)}{(2 - \xi - \eta)^2} UV + 2FV \right\} \, d\xi \, d\eta.
 \tag{3.17}$$

Now we integrate by parts in  $I_V$  in the following way:

- in the term  $U_\xi V_\eta$ , we move the derivative from  $V_\eta$  to  $U_\xi$  and obtain the term  $(U_\xi)_\eta V$ :

$$\int_D (\eta - \xi)^{2\beta} U_\xi V_\eta d\xi d\eta = - \int_D (\eta - \xi)^{2\beta} \left[ (U_\xi)_\eta + \frac{2\beta}{\eta - \xi} U_\xi \right] V d\xi d\eta; \tag{3.18}$$

- in the term  $U_\eta V_\xi$ , we move the derivative from  $U_\eta$  to  $V_\xi$  and obtain the term  $U(V_\xi)_\eta$ :

$$\int_D (\eta - \xi)^{2\beta} U_\eta V_\xi d\xi d\eta = - \int_D (\eta - \xi)^{2\beta} \left[ (V_\xi)_\eta + \frac{2\beta}{\eta - \xi} V_\xi \right] U d\xi d\eta.$$

There are not integrals on the boundary of  $D$  because  $U(0, \eta) = 0$ ,  $V(\xi, \eta) \equiv 0$  in a neighborhood of  $\{\eta = \xi\}$  and in a neighborhood of  $\{\eta = 1\}$ .

- since  $V \in C^2(\bar{D})$ , we have  $(V_\xi)_\eta = (V_\eta)_\xi$ ;
- in the term  $(V_\eta)_\xi U$ , we move the derivatives from  $(V_\eta)_\xi$  to  $U$  and obtain the term  $(U_\xi)_\eta V$ :

$$\begin{aligned} \int_D (\eta - \xi)^{2\beta} U_\eta V_\xi d\xi d\eta &= - \int_D (\eta - \xi)^{2\beta} \left[ (V_\eta)_\xi + \frac{2\beta}{\eta - \xi} V_\xi \right] U d\xi d\eta \\ &= \int_D (\eta - \xi)^{2\beta} \left[ U_\xi V_\eta - \frac{2\beta}{\eta - \xi} (V_\xi + V_\eta) U \right] d\xi d\eta \\ &= - \int_D (\eta - \xi)^{2\beta} \left[ (U_\xi)_\eta - \frac{2\beta}{\eta - \xi} U_\eta \right] V d\xi d\eta. \end{aligned} \tag{3.19}$$

Again there are not integrals on the boundary of  $D$ , and putting (3.18) and (3.19) into (3.17), we get

$$I_V = -2 \int_D (\eta - \xi)^{2\beta} \left\{ (U_\xi)_\eta + \frac{\beta}{\eta - \xi} (U_\xi - U_\eta) - \frac{n(n+1)}{(2 - \xi - \eta)^2} U - F \right\} V d\xi d\eta = 0. \tag{3.20}$$

(2.ii) Let  $V(\xi, \eta) \in V^{(2)}$  and  $\Psi(s)$  be a function having the properties  $\Psi(s) \in C^\infty(\mathbf{R}^1)$ ,  $\Psi(s) = 1$  for  $s \geq 2$ ,  $\Psi(s) = 0$  for  $s \leq 1$ . If  $k, l \in \mathbf{N}$ , then according to (2.i) (see (3.17) and (3.20)) for the functions

$$V_{k,l}(\xi, \eta) := V(\xi, \eta) \Psi(k[1 - \eta]) \Psi(l[\eta - \xi])$$

identity (2.9) holds. Therefore we have

$$\begin{aligned} 0 &= \int_D (\eta - \xi)^{2\beta} \left\{ U_\xi V_\eta + U_\eta V_\xi + \frac{2n(n+1)}{(2 - \xi - \eta)^2} UV + 2FV \right\} \\ &\quad \times \Psi(k[1 - \eta]) \Psi(l[\eta - \xi]) d\xi d\eta \\ &\quad + \int_D l(\eta - \xi)^{2\beta} \{ U_\xi - U_\eta \} \Psi(k[1 - \eta]) \Psi'(l[\eta - \xi]) V d\xi d\eta \\ &\quad - \int_D k(\eta - \xi)^{2\beta} U_\xi \Psi'(k[1 - \eta]) \Psi(l[\eta - \xi]) V d\xi d\eta \\ &=: I_{1,kl} + I_{2,kl} + I_{3,kl}. \end{aligned} \tag{3.21}$$

Obviously,  $I_{1,kl} \rightarrow I_V$ , as  $k, l \rightarrow \infty$ .



We know that  $V \equiv 0$  in a neighborhood of  $(0, 0)$  and  $\text{supp } \Psi'(l[\eta - \xi])$  is contained in  $\{1 \leq l[\eta - \xi] \leq 2\}$ . Using estimate (3.12) we find that on  $\text{supp } \Psi'(l[\eta - \xi])$  the functions

$$W_{k,l}(\xi, \eta) := l(\eta - \xi)^{2\beta} \{U_\xi - U_\eta\} \Psi(k[1 - \eta]) \Psi'(l[\eta - \xi]) V$$

satisfy the estimates

$$|W_{k,l}(\xi, \eta)| \leq \text{const.}(\eta - \xi)^{\beta-1}. \tag{3.22}$$

Since, obviously, the sequence  $W_{k,l}$  converges pointwise almost everywhere to zero and it is dominated by a Lebesgue integrable function in  $D$  for  $0 < \beta < 1$  (see (3.22)). Thus, according to the Lebesgue dominated convergence theorem,  $I_{2,kl} \rightarrow 0$  as  $k, l \rightarrow \infty$ .

Since  $V(\xi, 1) = 0$ , we have

$$k|V(\xi, \eta)| |\Psi'(k[1 - \eta])| = k(1 - \eta) |V_\eta(\xi, \tilde{\eta})| |\Psi'(k[1 - \eta])| \leq c_\nu,$$

where  $c_\nu$  is a constant and  $\eta < \tilde{\eta} < 1$ . Therefore  $I_{3,kl} \rightarrow 0$  as  $k, l \rightarrow \infty$ .

Thus, letting  $k, l \rightarrow \infty$  in (3.21), we obtain that identity (2.9) holds for  $V \in V^{(2)}$ . Consequently, the function  $U(\xi, \eta)$  is a generalized solution to problem  $PK_2$ . □

#### 4 Proof of the main results

In this section we give the proofs of Theorem 1.1, Theorem 1.2 and Theorem 1.3 formulated in Section 1.

*Proof of Theorem 1.1* Let  $u_1$  and  $u_2$  be two generalized solutions of problem  $PK$  in  $\Omega_m$ . Then the function  $u := u_1 - u_2$  solves the homogeneous problem  $PK$ . We will show that the Fourier expansion

$$u(r, \theta, \varphi, t) = \sum_{n=0}^{\infty} \sum_{s=1}^{2n+1} u_n^s(r, t) Y_n^s(\theta, \varphi)$$

has zero Fourier-coefficients

$$u_n^s(r, t) := \int_0^\pi \int_0^{2\pi} u(r, \theta, \varphi, t) Y_n^s(\theta, \varphi) \sin \theta \, d\varphi \, d\theta$$

in  $G_m$ , i.e.,  $u \equiv 0$  in  $\Omega_m$ .

For  $u$  we know that the identity

$$\int_{\Omega_m} \left\{ t^m u_t v_t - u_{x_1} v_{x_1} - u_{x_2} v_{x_2} - u_{x_3} v_{x_3} \right\} dx_1 \, dx_2 \, dx_3 \, dt = 0 \tag{4.1}$$

holds for all test functions  $v(x, t) = w(r, t) Y_n^s(x)$  described in Remark 2.1. Therefore from (4.1) we derive

$$\int_{G_m} \left\{ u_{n,r}^s w_r - t^m u_{n,t}^s w_t + \frac{n(n+1)}{r^2} u_n^s w \right\} r^2 \, dr \, dt = 0 \tag{4.2}$$

for all  $w(r, t) \in V_m^{(1)}$  (see Definition 2.1),  $n \in \mathbf{N} \cup \{0\}$ ,  $s = 1, 2, \dots, 2n + 1$ . Since  $u(x, t)$  satisfies conditions (1), (2) and (3) in Definition 1.1, the functions  $u_n^s(r, t)$  satisfy conditions (1), (2) and (3) in Definition 2.1, and therefore they are generalized solutions of problem  $PK_1$ .

Using (2.3) we see that the functions  $V(\xi, \eta) := r(\xi, \eta)w(r(\xi, \eta), t(\xi, \eta)) \in V^{(2)}$ . Now from (4.2) we obtain that for the functions  $U_n^s(\xi, \eta) := r(\xi, \eta)u_n^s(r(\xi, \eta), t(\xi, \eta))$  the identity

$$\int_D (\eta - \xi)^{2\beta} \left\{ U_{n,\xi}^s V_\eta + U_{n,\eta}^s V_\xi + \frac{2n(n+1)}{(2-\xi-\eta)^2} U_n^s V \right\} d\xi d\eta = 0$$

holds for all  $V(r, t) \in V^{(2)}$  (see Definition 2.2),  $n \in \mathbf{N} \cup \{0\}$ ,  $s = 1, 2, \dots, 2n + 1$ . The functions  $U_n^s(\xi, \eta)$  satisfy conditions (1), (2) and (3) in Definition 2.2 and, consequently,  $U_n^s(\xi, \eta)$  are generalized solutions of the 2-D homogeneous problem  $PK_2$ . Theorem 3.1 gives  $U_n^s(\xi, \eta) \equiv 0$  in  $D$ . Therefore  $u_n^s(r, t) \equiv 0$  in  $G_m$  and thus  $u = u_1 - u_2 \equiv 0$  in  $\Omega_m$ .  $\square$

*Proof of Theorem 1.2* From Theorem 1.1 it follows that there exists at most one generalized solution of problem  $PK$  in  $\Omega_m$ . Since  $f(x, t)$  has the form (1.5), we look for a generalized solution of the form (1.6), i.e.,

$$u(x, t) = \sum_{n=0}^l \sum_{s=1}^{2n+1} u_n^s(|x|, t) Y_n^s(x).$$

To find such a solution means to find functions  $u_n^s(r, t)$  that satisfy the identities

$$\int_{G_{m,s}} \left[ u_{n,r}^s v_r - t^m u_{n,t}^s v_t + \frac{n(n+1)}{r^2} u_n^s v + f_n^s v \right] r^2 dr dt = 0$$

for all  $v \in V_m^{(1)}$  and satisfy the corresponding conditions (1), (2) and (3) in Definition 2.1. In view of (2.3) to find such functions means to find functions

$$U_n^s(\xi, \eta) = r(\xi, \eta)u_n^s(r(\xi, \eta), t(\xi, \eta)),$$

such that for  $F_n^s(\xi, \eta) := \frac{1}{4}r(\xi, \eta)f_n^s(r(\xi, \eta), t(\xi, \eta))$  the identity

$$\int_D (\eta - \xi)^{2\beta} \left\{ U_{n,\xi}^s V_\eta + U_{n,\eta}^s V_\xi + \frac{2n(n+1)}{(2-\xi-\eta)^2} U_n^s V + 2F_n^s V \right\} d\xi d\eta = 0$$

holds for all  $V(\xi, \eta) = r(\xi, \eta)v(r(\xi, \eta), t(\xi, \eta)) \in V^{(2)}$  and satisfies the corresponding conditions (1), (2) and (3) in Definition 2.2. Theorem 3.5 gives the existence of such functions  $U_n^s(\xi, \eta)$  which are generalized solutions of problem  $PK_2$  in  $D$ . In that way we found functions  $u_n^r(r, t) = r^{-1}U_n^s(\xi(r, t), \eta(r, t))$  which are generalized solutions of problem  $PK_1$  in  $G_m$ . Therefore the function  $u(x, t)$ , given by (1.6), is a generalized solution of problem  $PK$  in  $\Omega_m$ .  $\square$

*Proof of Theorem 1.3* Theorem 1.1 and Theorem 1.2 claim the existence and uniqueness of generalized solutions  $u(x, t)$  of problem  $PK$  in  $\Omega_m$ , which has the form (1.6). Using (2.3) for functions  $U_n^s(\xi, \eta) = r(\xi, \eta)u_n^s(r(\xi, \eta), t(\xi, \eta))$  and  $F_n^s(\xi, \eta) = \frac{1}{4}r(\xi, \eta)f_n^s(r(\xi, \eta), t(\xi, \eta))$ , we obtain the 2-D problem  $PK_2$ . According to Theorem 3.5, estimates (3.12) hold, and in view

of (3.6) and (3.7) we see that the estimate for  $|U_n^s(\xi, \eta)|$  holds with  $(\max_{\bar{G}_m} |f_n^s|)$  instead of  $M_F$ :

$$|U_n^s(\xi, \eta)| \leq K \left( \max_{\bar{G}_m} |f_n^s| \right) (2 - \xi - \eta)^{-n}$$

with a constant  $K > 0$  independent of  $f_n^s$ . That implies

$$|u_n^s(r, t)| \leq 2^{-n} K \left( \max_{\bar{G}_m} |f_n^s| \right) r^{-n-1}.$$

Therefore in view of (1.6), summing up over  $n$  and  $s$ , we get the desired estimate (1.7).  $\square$

**Remark 4.1** It is interesting that in the case  $0 < m < 1$  problem  $PK$  for the Keldysh-type equation (1.1) can be formally reduced to problem  $P2$  for the Tricomi-type equation (1.2) with power of degeneration  $m_1 := m/(1 - m) > 0$  and the right-hand side function, which vanishes on  $\Sigma_0$  like  $t^{m_1}$ . That implies many differences between the investigation and behavior of the solution to the obtained problem and the usual Protter problem  $P2$ . However, in the present paper we study the  $(3 + 1)$ -D problem  $PK$  in a more general case when  $0 < m < 4/3$ .

**Appendix A: The Riemann-Hadamard function  $\Phi(\xi, \eta, \xi_0, \eta_0)$**

Firstly, to aid the reader, we briefly recall some known properties of the hypergeometric function of Gauss  $F(a, b, c; \zeta)$  that we will use.

If  $c \neq 0, -1, -2, \dots$ , then

$$F(a, b, c; \zeta) := \sum_{i=0}^{\infty} \frac{(a)_i (b)_i}{i! (c)_i} \zeta^i, \tag{A.1}$$

with  $(a)_i = \Gamma(a + i)/\Gamma(a)$ , where  $\Gamma$  is the Euler gamma function of Euler. For  $i \in \mathbf{N}$ , one has  $(a)_i = a(a + 1) \cdots (a + i - 1)$ ,  $(a)_0 = 1$ .

The series (A.1) converges absolutely for  $\zeta \in \mathbf{C}$  with  $|\zeta| < 1$  and also for  $|\zeta| = 1$  if  $\text{Re}(c - a - b) > 0$ . If  $-1 < \text{Re}(c - a - b) < 0$ , then the series converges conditionally for  $|\zeta| = 1$  with  $\zeta \neq 1$ .

We mention the following properties of the hypergeometric function (see [31, 49, 50]):

$$F(a, b, c; \zeta) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c - a)} \int_0^1 t^{a-1} (1 - t)^{c-a-1} (1 - \zeta t)^{-b} dt \tag{A.2}$$

for  $\zeta \in \mathbf{C}$ ,  $0 < \text{Re}(a) < \text{Re}(c)$ ,  $|\arg(1 - \zeta)| < \pi$ .

In the case  $c - a - b > 0$ :

$$|F(a, b, c; \zeta)| \leq \text{const.}, \quad F(a, b, c; 1) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}; \tag{A.3}$$

resp.  $c - a - b < 0$ :

$$F(a, b, c; \zeta) = (1 - \zeta)^{c-a-b} F(c - a, c - b, c; \zeta) \tag{A.4}$$

and

$$|F(a, b, c; \zeta)| \leq \text{const.}(1 - \zeta)^{c-a-b}; \tag{A.5}$$

resp.  $c - a - b = 0$ : For each  $\alpha > 0$ , there exists a constant  $c(\alpha) > 0$  such that

$$|F(a, b, c; \zeta)| \leq c(\alpha)(1 - \zeta)^{-\alpha}, \tag{A.6}$$

$$\frac{d}{d\zeta}F(a, b, c; \zeta) = \frac{ab}{c}F(a + 1, b + 1, c + 1; \zeta). \tag{A.7}$$

The hypergeometric function of two variables is defined by

$$F_1(a, b_1, b_2, c; x, y) := \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{(a)_{i+j}(b_1)_j(b_2)_i}{(c)_{i+j}i!j!} x^j y^i. \tag{A.8}$$

The series converges absolutely for  $x, y \in \mathbf{C}$  with  $|x| < 1, |y| < 1$  (for more properties of  $F_1$ , see [49], pp.224-228).

Now, in the case  $n \in \mathbf{N} \cup \{0\}$ , we construct the following Riemann-Hadamard function of the form (3.1) associated to problem  $PK_2$ : For  $(\xi_0, \eta_0) \in D$

$$\begin{aligned} \Phi^+ &= \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta F_3(\beta, n + 1, 1 - \beta, -n, 1; X, Y), \quad \eta > \xi_0, \\ \Phi^- &= k \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta X^{-\beta} H_2\left(\beta, \beta, -n, n + 1, 2\beta; \frac{1}{X}, -Y\right), \quad \eta < \xi_0, \end{aligned} \tag{A.9}$$

where

$$\begin{aligned} k &= \frac{\Gamma(\beta)}{\Gamma(1 - \beta)\Gamma(2\beta)}, \\ X &= X(\xi, \eta, \xi_0, \eta_0) := \frac{(\xi_0 - \xi)(\eta_0 - \eta)}{(\eta - \xi)(\eta_0 - \xi_0)}, \\ Y &= Y(\xi, \eta, \xi_0, \eta_0) := -\frac{(\xi_0 - \xi)(\eta_0 - \eta)}{(2 - \xi - \eta)(2 - \xi_0 - \eta_0)}. \end{aligned}$$

Here  $F_3(a_1, a_2, b_1, b_2, c; x, y)$  is the Appell series

$$F_3(a_1, a_2, b_1, b_2, c; x, y) := \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{(a_1)_j(a_2)_i(b_1)_j(b_2)_i}{(c)_{i+j}i!j!} x^j y^i \tag{A.10}$$

which converges absolutely for  $x, y \in \mathbf{C}$  with  $|x| < 1, |y| < 1$  (see [49], pp.224-228) and  $H_2(a_1, a_2, b_1, b_2, c; x, y)$  is the Horn series

$$H_2(a_1, a_2, b_1, b_2, c; x, y) := \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{(a_1)_{j-i}(a_2)_j(b_1)_i(b_2)_i}{(c)_j i! j!} x^j y^i \tag{A.11}$$

which converges absolutely for  $x, y \in \mathbf{C}$  with  $|x| < 1, |y|(1 + |x|) < 1$  (see [49], pp.224-228).

We mention that for  $(\xi_0, \eta_0) \in D$  we have  $|X| < 1$  in  $\bar{\Pi}$  and  $1/|X| < 1$  in  $\bar{T}$ , while  $|Y| < 1$  in  $\bar{\Pi}$  but  $|Y|$  could be greater than 1 in  $T$ . However, the function  $\Phi$  is well defined because

$n \in \mathbf{N} \cup \{0\}$ , since  $b_1 = -n$ , and we have a finite sum with respect to  $i$  in the function  $H_2$  (see (A.11)), which appears in (A.9). We will fix all these properties a little bit later.

Let, for  $(\xi_0, \eta_0) \in D$  and  $(\xi, \eta) \in \bar{T} \cup \bar{\Pi} \setminus \{\eta = \xi_0\}$ , us introduce the functions

$$H(\xi, \eta; \xi_0, \eta_0) = \begin{cases} H^+(\xi, \eta; \xi_0, \eta_0), & \eta > \xi_0, \\ H^-(\xi, \eta; \xi_0, \eta_0), & \eta < \xi_0, \end{cases} \tag{A.12}$$

where

$$H^+(\xi, \eta; \xi_0, \eta_0) = \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta F(\beta, 1 - \beta, 1; X),$$

$$H^-(\xi, \eta; \xi_0, \eta_0) = k \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta X^{-\beta} F\left(\beta, \beta, 2\beta; \frac{1}{X}\right)$$

and

$$G(\xi, \eta; \xi_0, \eta_0) = \begin{cases} G^+(\xi, \eta; \xi_0, \eta_0), & \eta > \xi_0, \\ G^-(\xi, \eta; \xi_0, \eta_0), & \eta < \xi_0, \end{cases} \tag{A.13}$$

where

$$G^+(\xi, \eta; \xi_0, \eta_0) := \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta \sum_{i=1}^n c_i Y^i F(\beta, 1 - \beta, i + 1; X), \tag{A.14}$$

$$G^-(\xi, \eta; \xi_0, \eta_0) := k \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta X^{-\beta} \sum_{i=1}^n d_i Y^i F\left(\beta - i, \beta, 2\beta; \frac{1}{X}\right), \tag{A.15}$$

$$c_i := \frac{(n + 1)_i (-n)_i}{i! i!}, \quad d_i := \frac{(n + 1)_i (-n)_i}{(1 - \beta)_i i!}.$$

Now we prove the following important lemma.

**Lemma A.1** *The function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  has the following decomposition:*

$$\Phi(\xi, \eta; \xi_0, \eta_0) = H(\xi, \eta; \xi_0, \eta_0) + G(\xi, \eta; \xi_0, \eta_0). \tag{A.16}$$

*Proof* (i) In view of (A.10) we have

$$F_3(\beta, n + 1, 1 - \beta, -n, 1; X, Y) = \sum_{i=0}^n \sum_{j=0}^\infty \frac{(\beta)_j (1 - \beta)_j (n + 1)_i (-n)_i}{(1)_{i+j} i! j!} X^i Y^j.$$

Since  $(1)_{i+j} = (i + j)! = i!(i + 1)_j$  for  $i, j \in \mathbf{N} \cup \{0\}$ , we obtain from (A.1) and (A.9)

$$\begin{aligned} \Phi^+(\xi, \eta; \xi_0, \eta_0) &= \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta \left\{ \sum_{j=0}^\infty \frac{(\beta)_j (1 - \beta)_j}{(1)_{jj}!} X^j + \sum_{i=1}^n c_i Y^i \sum_{j=0}^\infty \frac{(\beta)_j (1 - \beta)_j}{(i + 1)_{jj}!} X^j \right\} \\ &= \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta \left\{ F(\beta, 1 - \beta, 1; X) + \sum_{i=1}^n c_i Y^i F(\beta, 1 - \beta, i + 1; X) \right\} \\ &= H^+(\xi, \eta; \xi_0, \eta_0) + G^+(\xi, \eta; \xi_0, \eta_0). \end{aligned}$$

(ii) In view of (A.11) we have

$$H_2\left(\beta, \beta, -n, n + 1, 2\beta; \frac{1}{X}, Y\right) := \sum_{i=0}^n \sum_{j=0}^{\infty} \frac{(\beta)_{j-i}(\beta)_j(-n)_i(1-n)_i}{(2\beta)_{j!}} X^{-j}(-Y)^i.$$

We mention that for  $0 < \beta < 1$  and  $i, j \in \mathbf{N} \cup \{0\}$

$$(\beta)_{j-i} = \frac{\Gamma(\beta + j - i)}{\Gamma(\beta)} = \frac{\Gamma(\beta - i)}{\Gamma(\beta)}(\beta - i)_j,$$

and using the relation  $\Gamma(z)\Gamma(1 - z) = \frac{\pi}{\sin(\pi z)}$  we calculate

$$\begin{aligned} (\beta)_{j-i}(1 - \beta)_i &= (\beta - i)_j \frac{\Gamma(\beta - i)\Gamma(1 - \beta + i)}{\Gamma(\beta)\Gamma(1 - \beta)} \\ &= (\beta - i)_j \frac{\sin(\beta\pi)}{\sin((\beta - i)\pi)} \\ &= (-1)^i(\beta - i)_j. \end{aligned}$$

Now, from (A.1) and (A.9) we obtain

$$\begin{aligned} \Phi^-(\xi, \eta; \xi_0, \eta_0) &= k \left( \frac{\eta - \xi}{\eta_0 - \xi_0} \right)^\beta X^{-\beta} \left\{ \sum_{j=0}^{\infty} \frac{(\beta)_j(\beta)_j}{(2\beta)_{j!}} \frac{1}{X^j} + \sum_{i=1}^n d_i Y^i \sum_{j=0}^{\infty} \frac{(\beta - i)_j(\beta)_j}{(2\beta)_{j!}} \frac{1}{X^j} \right\} \\ &= k \left( \frac{\eta - \xi}{\eta_0 - \xi_0} \right)^\beta X^{-\beta} \left\{ F\left(\beta, \beta, 2\beta; \frac{1}{X}\right) + \sum_{i=1}^n d_i Y^i F\left(\beta - i, \beta, 2\beta; \frac{1}{X}\right) \right\} \\ &= H^-(\xi, \eta; \xi_0, \eta_0) + G^-(\xi, \eta; \xi_0, \eta_0). \quad \square \end{aligned}$$

We mention here that function (A.9) is closely connected to the Riemann-Hadamard function announced in [51], p.25, example 7, which is associated to a Cauchy-Goursat problem for an equation connected with (2.5) with some appropriate substitutions. Actually, the function  $H(\xi, \eta; \xi_0, \eta_0)$  is the Riemann-Hadamard function associated to problem  $PK_2$  in the case  $n = 0$  (see Gellerstedt [46] and Smirnov [47]). It is well known that the function  $H(\xi, \eta; \xi_0, \eta_0)$  has the properties (i) ÷ (vi) listed in Section 3. It is not difficult to check that in the case  $n \geq 0$  function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  has the same properties. Using the systems of differential equations that  $F_3$  and  $H_2$  satisfy (see [49], pp.233-234), with a straightforward calculation we check that the function  $\Phi(\xi, \eta; \xi_0, \eta_0)$  satisfies equations (3.2) and (3.3). Further, since  $X(\xi_0, \eta, \xi_0, \eta_0) = X(\xi, \eta_0, \xi_0, \eta_0) = 0$ ,  $Y(\xi_0, \eta, \xi_0, \eta_0) = Y(\xi, \eta_0, \xi_0, \eta_0) = 0$ , we see that the function  $\Phi$  has the properties (ii), (iii) and (iv). We also have  $X(\xi, \xi, \xi_0, \eta_0) = 0$ , and therefore the function  $G(\xi, \eta; \xi_0, \eta_0)$  vanishes on the line  $\{\eta = \xi\}$  of power  $2\beta$ . Therefore the function  $\Phi$  has the properties (vi). Let us calculate the jump of the function  $\Phi$  on the line  $\{\eta = \xi_0\}$ . We will show that the function  $G$  has no jump on the line  $\{\eta = \xi_0\}$ . Using (A.3) and the relation  $\Gamma(i) = (i - 1)!$  for  $i \in \mathbf{N}$ , we calculate

$$c_i F(\beta, 1 - \beta, i + 1; 1) = k d_i F(\beta - i, \beta, 2\beta; 1) = \frac{(n + 1)_i(-n)_i}{i\Gamma(1 - \beta + i)\Gamma(\beta + i)}.$$

In view of (A.14) and (A.15) we have

$$G^+(\xi, \xi_0; \xi_0, \eta_0) = G^-(\xi, \xi_0; \xi_0, \eta_0) = \left(\frac{\xi_0 - \xi}{\eta_0 - \xi_0}\right)^\beta \sum_{i=1}^n \frac{(n+1)_i (-n)_i}{i \Gamma(1-\beta+i) \Gamma(\beta+i)} Y^i(\xi, \xi_0, \xi_0, \eta_0).$$

Therefore the jump  $[[G]] = 0$ , and in view of (A.16) we have  $[[\Phi]] = [[H]]$ . Consequently, the function  $\Phi$  has the property (v) since  $[[H]] = \cos(\pi\beta) \left(\frac{\xi_0 - \xi}{\eta_0 - \xi_0}\right)^\beta$  (see Gellerstedt [46]).

### A.1 The function $H(\xi, \eta, \xi_0, \eta_0)$

Using the properties of a hypergeometric function mentioned above and the relations

$$1 - X = \frac{(\eta_0 - \xi)(\eta - \xi_0)}{(\eta - \xi)(\eta_0 - \xi_0)}, \quad 1 - \frac{1}{X} = \frac{(\eta_0 - \xi)(\xi_0 - \eta)}{(\xi_0 - \xi)(\eta_0 - \eta)},$$

$$X_{\xi_0} = \frac{(\eta_0 - \xi)(\eta_0 - \eta)}{(\eta - \xi)(\eta_0 - \xi_0)^2}, \quad X_{\eta_0} = \frac{(\eta - \xi_0)(\xi_0 - \xi)}{(\eta - \xi)(\eta_0 - \xi_0)^2}, \tag{A.17}$$

$$\left(\frac{1}{X}\right)_{\xi_0} = \frac{(\xi - \eta_0)(\eta - \xi)}{(\eta_0 - \eta)(\xi_0 - \xi)^2}, \quad \left(\frac{1}{X}\right)_{\eta_0} = \frac{(\xi_0 - \eta)(\eta - \xi)}{(\xi_0 - \xi)(\eta_0 - \eta)^2}, \tag{A.18}$$

we prove the following lemma.

**Lemma A.2** *Let  $0 < \beta < 1$  and  $0 < \xi_0 < \eta_0 < 1$ . Then there exists a constant  $C_H > 0$  such that*

$$|H^+(\xi, \eta; \xi_0, \eta_0)| \leq C_H (\eta - \xi_0)^{-\beta}, \quad (\xi, \eta) \in \Pi, \tag{A.19}$$

$$|H^-(\xi, \eta; \xi_0, \eta_0)| \leq C_H (\xi_0 - \eta)^{-\beta}, \quad (\xi, \eta) \in T, \tag{A.20}$$

$$|H_{\eta_0}^+(\xi, \eta; \xi_0, \eta_0)| \leq C_H \frac{(\eta - \xi_0)^{-\beta}}{\eta_0 - \xi_0}, \quad (\xi, \eta) \in \Pi, \tag{A.21}$$

$$|H_{\eta_0}^-(\xi, \eta; \xi_0, \eta_0)| \leq C_H \frac{(\xi_0 - \eta)^{-\beta}}{\eta_0 - \eta}, \quad (\xi, \eta) \in T. \tag{A.22}$$

*Proof* (i) Using (A.6) we find that for each  $\alpha > 0$  there exists a constant  $c(\alpha) > 0$  such that

$$|H^+(\xi, \eta; \xi_0, \eta_0)| \leq c(\alpha) \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta (1 - X)^{-\alpha} = c(\alpha) \frac{(\eta - \xi)^{\alpha+\beta} (\eta_0 - \xi_0)^{\alpha-\beta}}{(\eta - \xi_0)^\alpha (\eta_0 - \xi)^\alpha},$$

$$|H^-(\xi, \eta; \xi_0, \eta_0)| \leq c(\alpha) \frac{(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta (\eta_0 - \eta)^\beta} \left(1 - \frac{1}{X}\right)^{-\alpha} = c(\alpha) \frac{(\eta - \xi)^{2\beta} (\xi_0 - \xi)^{\alpha-\beta} (\eta_0 - \eta)^{\alpha-\beta}}{(\eta_0 - \xi)^\alpha (\xi_0 - \eta)^\alpha}.$$
(A.23)

From here, choosing  $\alpha = \beta$ , we obtain estimates (A.19), (A.20).

(ii) In view of (A.17), (A.18) for the derivatives with respect to  $\eta_0$ , using (A.4) and (A.7), we obtain

$$\begin{aligned} |H_{\eta_0}^+| &= \left| -\frac{\beta}{\eta_0 - \xi_0} H^+ + \beta(1 - \beta) \left( \frac{\eta - \xi}{\eta_0 - \xi_0} \right)^\beta X_{\eta_0} F(1 + \beta, 2 - \beta, 2; X) \right| \\ &= \frac{\beta(\eta - \xi)^\beta}{(\eta_0 - \xi_0)^{1+\beta}} \left| -F(\beta, 1 - \beta, 1; X) + (1 - \beta) \frac{\xi_0 - \xi}{\eta_0 - \xi} F(1 - \beta, \beta, 2; X) \right| \\ &\leq c(\alpha) \frac{(\eta - \xi)^\beta}{(\eta_0 - \xi_0)^{1+\beta}} (1 - X)^{-\alpha} \leq c(\alpha) \frac{(\eta_0 - \xi_0)^{\alpha-\beta-1}}{(\eta - \xi_0)^\alpha}, \end{aligned}$$

and

$$\begin{aligned} |H_{\eta_0}^-| &= \left| \frac{\beta H^-}{\eta - \eta_0} + \frac{k\beta(\eta - \xi)^{2\beta}}{2(\xi_0 - \xi)^\beta(\eta_0 - \eta)^\beta} \left( \frac{1}{X} \right)_{\eta_0} F\left(1 + \beta, 1 + \beta, 1 + 2\beta; \frac{1}{X}\right) \right| \\ &= \frac{k\beta(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta(\eta_0 - \eta)^{1+\beta}} \left| -F\left(\beta, \beta, 2\beta; \frac{1}{X}\right) + \frac{1}{2} \frac{\eta - \xi}{\eta_0 - \xi} F\left(\beta, \beta, 1 + 2\beta; \frac{1}{X}\right) \right| \\ &\leq c(\alpha) \frac{(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta(\eta_0 - \eta)^{1+\beta}} \left(1 - \frac{1}{X}\right)^{-\alpha} \leq c(\alpha) \frac{(\eta_0 - \eta)^{\alpha-\beta-1}}{(\xi_0 - \eta)^\alpha}. \end{aligned}$$

Now we choose  $\alpha = \beta$  to obtain the desired estimates (A.21), (A.22). □

**Remark A.1** In the same manner, for the derivatives with respect to  $\xi_0$ , we obtain

$$H_{\xi_0}^+ = \beta \frac{(\eta - \xi)^\beta}{(\eta_0 - \xi_0)^{1+\beta}} \left[ F(\beta, 1 - \beta, 1; X) + (1 - \beta) \frac{\eta_0 - \eta}{\eta - \xi_0} F(1 - \beta, \beta, 2; X) \right] \tag{A.24}$$

and

$$H_{\xi_0}^- = -\frac{k\beta(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^{1+\beta}(\eta_0 - \eta)^\beta} \left[ F\left(\beta, \beta, 2\beta; \frac{1}{X}\right) + \frac{1}{2} \frac{\eta - \xi}{\xi_0 - \eta} F\left(\beta, \beta, 1 + 2\beta; \frac{1}{X}\right) \right]. \tag{A.25}$$

In the case  $0 < \beta < 1/2$ , Smirnov [47] and Meredov [52] claim

$$|H_{\xi_0}^+(\xi, \eta; \xi_0, \eta_0)| \leq \frac{(\eta - \xi)(\eta_0 - \xi_0)^{-2\beta}}{(\eta_0 - \xi)^{1-\beta}(\eta - \xi_0)^{1-\beta}},$$

i.e.,  $H_{\xi_0}^+(\xi, \eta; \xi_0, \eta_0)$  has integrable singularity on  $\{\eta = \xi_0\}$ . As we see from (A.24) and (A.25), the derivative with respect to  $\xi_0$  of function  $H$  has not integrable singularity on  $\{\eta = \xi_0\}$ .

### A.2 The function $G(\xi, \eta, \xi_0, \eta_0)$

In this section we prove some properties of the function  $G(\xi, \eta; \xi_0, \eta_0)$  defined by (A.13).

**Lemma A.3** *Let  $0 < \beta < 1$  and  $0 < \xi_0 < \eta_0 < 1$ . Then there exists a constant  $C_G > 0$  such that*

$$|G^+(\xi, \eta; \xi_0, \eta_0)| \leq C_G(\eta_0 - \xi_0)^{-\beta}, \quad (\xi, \eta) \in \Pi, \tag{A.26}$$

$$|G^-(\xi, \eta; \xi_0, \eta_0)| \leq C_G(2 - \xi_0 - \eta_0)^{-n}, \quad (\xi, \eta) \in T, \tag{A.27}$$



$$|G_{\eta_0}^+(\xi, \eta, \xi_0, \eta_0)| \leq C_G \frac{(\eta_0 - \xi_0)^{-\beta}}{2 - \xi_0 - \eta_0}, \quad (\xi, \eta) \in \Pi, \tag{A.28}$$

$$|G_{\eta_0}^-(\xi, \eta, \xi_0, \eta_0)| \leq C_G \frac{(\eta_0 - \eta)^{-\beta}}{(2 - \xi_0 - \eta_0)^{n+1}}, \quad (\xi, \eta) \in T, \tag{A.29}$$

$$|G_{\xi_0}^+(\xi, \eta, \xi_0, \eta_0)| \leq C_G \frac{(\eta - \xi_0)^{-\beta}}{(2 - \xi_0 - \eta_0)}, \quad (\xi, \eta) \in \Pi, \tag{A.30}$$

$$|G_{\xi_0}^-(\xi, \eta, \xi_0, \eta_0)| \leq C_G \frac{(\xi_0 - \eta)^{-\beta}}{(2 - \xi_0 - \eta_0)^{n+1}}, \quad (\xi, \eta) \in T. \tag{A.31}$$

*Proof* First, we mention that

$$Y_{\xi_0} = -\frac{(2 - \xi - \eta_0)(\eta_0 - \eta)}{(2 - \xi - \eta)(2 - \xi_0 - \eta_0)^2}, \quad Y_{\eta_0} = -\frac{(2 - \xi_0 - \eta)(\xi_0 - \xi)}{(2 - \xi - \eta)(2 - \xi_0 - \eta_0)^2}.$$

(i) Let  $(\xi, \eta) \in \Pi$ . Then we have

$$\begin{aligned} |X_{\xi_0}| &\leq \frac{\eta_0 - \xi}{(\eta - \xi)(\eta_0 - \xi_0)}, \\ |Y| < 1, \quad \frac{|Y|}{\eta_0 - \xi_0} &\leq \frac{|Y|(\eta_0 - \xi)}{(\eta - \xi)(\eta_0 - \xi_0)} \leq \frac{1}{2 - \xi_0 - \eta_0}, \\ |Y_{\xi_0}| &\leq \frac{1}{2 - \xi_0 - \eta_0}, \quad |Y_{\eta_0}| \leq \frac{2}{2 - \xi_0 - \eta_0}. \end{aligned} \tag{A.32}$$

According to (A.3),  $|F(\beta, 1 - \beta, i + 1; X)| \leq const., i = 1, 2, \dots, n$ , in expression (A.14) for  $G^+$ . Therefore estimate (A.26) holds.

With use of (A.7) we calculate the derivative with respect to  $\xi_0$

$$\begin{aligned} G_{\xi_0}^+ &= \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta \left\{ \sum_{i=1}^n c_i \left[ \frac{\beta Y^i}{\eta_0 - \xi_0} + i Y^{i-1} Y_{\xi_0} \right] F(\beta, 1 - \beta, i + 1; X) \right. \\ &\quad \left. + \beta(1 - \beta) \sum_{i=1}^n \frac{c_i}{i + 1} Y^i X_{\xi_0} F(\beta + 1, 2 - \beta, i + 2; X) \right\} \end{aligned}$$

and the derivative with respect to  $\eta_0$

$$\begin{aligned} G_{\eta_0}^+ &= \left(\frac{\eta - \xi}{\eta_0 - \xi_0}\right)^\beta \left\{ \sum_{i=1}^n c_i \left[ -\frac{\beta Y^i}{\eta_0 - \xi_0} + i Y^{i-1} Y_{\eta_0} \right] F(\beta, 1 - \beta, i + 1; X) \right. \\ &\quad \left. + \beta(1 - \beta) \sum_{i=1}^n \frac{c_i}{i + 1} Y^i X_{\eta_0} F(\beta + 1, 2 - \beta, i + 2; X) \right\}. \end{aligned}$$

According to (A.3) and (A.6), for the hypergeometric functions in the expressions for  $G_{\xi_0}^+$  and  $G_{\eta_0}^+$ , we have

$$\begin{aligned} |F(\beta + 1, 2 - \beta, 3; X)| &\leq c(\alpha) \frac{(\eta_0 - \xi_0)^\alpha (\eta - \xi)^\alpha}{(\eta - \xi_0)^\alpha (\eta_0 - \xi)^\alpha} \leq c(\alpha) \left(\frac{\eta_0 - \xi_0}{\eta - \xi_0}\right)^\alpha, \quad \alpha > 0, \\ |F(1 + \beta, 2 - \beta, i + 2; X)| &\leq const., \quad i = 2, 3, \dots, n. \end{aligned}$$

Therefore in view of (A.17) we have

$$\begin{aligned}
 |G_{\xi_0}^+| &\leq \frac{C_1(\alpha)}{(\eta_0 - \xi_0)^\beta} \left\{ \frac{|Y|}{\eta_0 - \xi_0} + \frac{1}{2 - \xi_0 - \eta_0} + \frac{|Y|(\eta_0 - \xi)}{(\eta - \xi)(\eta_0 - \xi_0)} \left( \frac{\eta_0 - \xi_0}{\eta - \xi_0} \right)^\alpha \right\}, \\
 |G_{\eta_0}^+| &\leq \frac{C_2(\alpha)}{(\eta_0 - \xi_0)^\beta} \left\{ \frac{|Y|}{\eta_0 - \xi_0} + \frac{2}{2 - \xi_0 - \eta_0} + \frac{|Y|(\xi_0 - \xi)}{(\eta - \xi)(\eta_0 - \xi_0)} \left( \frac{\eta - \xi_0}{\eta_0 - \xi_0} \right)^{1-\alpha} \right\}.
 \end{aligned}$$

Now, taking  $\alpha = \beta \in (0, 1)$  and using (A.32), we obtain estimates (A.30) and (A.28).

(ii) Let  $(\xi, \eta) \in T$ . Then we have

$$\begin{aligned}
 \left| \left( \frac{1}{X} \right)_{\xi_0} \right| &\leq \frac{\eta_0 - \xi}{(\xi_0 - \xi)(\eta_0 - \eta)}, \\
 |Y| &< \frac{1}{2 - \xi_0 - \eta_0}, \quad \frac{|Y|}{\eta_0 - \eta} \leq \frac{1}{2 - \xi_0 - \eta_0}, \\
 \frac{|Y|}{\xi_0 - \xi} &\leq \frac{|Y|(\eta_0 - \xi)}{(\xi_0 - \xi)(\eta_0 - \eta)} \leq \frac{1}{2 - \xi_0 - \eta_0}, \\
 |Y_{\xi_0}| &\leq \frac{1}{(2 - \xi_0 - \eta_0)^2}, \quad |Y_{\eta_0}| \leq \frac{1}{(2 - \xi_0 - \eta_0)^2}.
 \end{aligned} \tag{A.33}$$

According to (A.3)  $|F(\beta - i, \beta, 2\beta; 1/X)| \leq const., i = 1, 2, \dots, n$ , in expression (A.15) for  $G^-$ . Since

$$\left( \frac{\eta - \xi}{\eta_0 - \xi_0} \right)^\beta X^{-\beta} = \frac{(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta (\eta_0 - \eta)^\beta},$$

we see that estimate (A.27) holds.

With use of (A.7) we calculate the derivative with respect to  $\xi_0$

$$\begin{aligned}
 G_{\xi_0}^- &= \frac{k(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta (\eta_0 - \eta)^\beta} \left\{ \sum_{i=1}^n d_i \left[ -\frac{\beta Y^i}{\xi_0 - \xi} + i Y^{i-1} Y_{\xi_0} \right] F\left(\beta - i, \beta, 2\beta; \frac{1}{X}\right) \right. \\
 &\quad \left. + \frac{1}{2} \sum_{i=1}^n (\beta - i) d_i Y^i \left( \frac{1}{X} \right)_{\xi_0} F\left(\beta - i + 1, \beta + 1, 2\beta + 1; \frac{1}{X}\right) \right\}
 \end{aligned}$$

and the derivative with respect to  $\eta_0$

$$\begin{aligned}
 G_{\eta_0}^- &= \frac{k(\eta - \xi)^{2\beta}}{(\xi_0 - \xi)^\beta (\eta_0 - \eta)^\beta} \left\{ \sum_{i=1}^n d_i \left[ -\frac{\beta Y^i}{\eta_0 - \eta} + i Y^{i-1} Y_{\eta_0} \right] F\left(\beta - i, \beta, 2\beta; \frac{1}{X}\right) \right. \\
 &\quad \left. + \frac{1}{2} \sum_{i=1}^n (\beta - i) d_i Y^i \left( \frac{1}{X} \right)_{\eta_0} F\left(\beta - i + 1, \beta + 1, 2\beta + 1; \frac{1}{X}\right) \right\}.
 \end{aligned}$$

According to (A.3) and (A.6), for the hypergeometric functions in the expressions for  $G_{\xi_0}^-$  and  $G_{\eta_0}^-$ , we have

$$\begin{aligned}
 \left| F\left(\beta, \beta + 1, 2\beta + 1; \frac{1}{X}\right) \right| &\leq c(\alpha) \frac{(\eta_0 - \eta)^\alpha (\xi_0 - \xi)^\alpha}{(\xi_0 - \eta)^\alpha (\eta_0 - \xi)^\alpha} \leq c(\alpha) \left( \frac{\eta_0 - \eta}{\xi_0 - \eta} \right)^\alpha, \quad \alpha > 0, \\
 |F(\beta - i + 1, 1 + \beta, 1 + 2\beta; 1/X)| &\leq const., \quad i = 2, 3, \dots, n.
 \end{aligned}$$

Now using (A.18) we calculate

$$\begin{aligned} |G_{\xi_0}^-| &\leq \frac{C_3(\alpha)}{(\eta_0 - \eta)^\beta} \left\{ \frac{|Y|}{\xi_0 - \xi} + \frac{1}{(2 - \xi_0 - \eta_0)^2} + \frac{|Y|(\eta_0 - \xi)}{(\xi_0 - \xi)(\eta_0 - \eta)} \left( \frac{\eta_0 - \eta}{\xi_0 - \eta} \right)^\alpha \right\} \sum_{i=1}^n |Y|^{i-1} \\ &\leq 3C_3(\alpha) \frac{(\eta_0 - \eta)^{\alpha-\beta}}{(\xi_0 - \eta)^\alpha} \sum_{i=1}^n (2 - \xi_0 - \eta_0)^{-i-1} \end{aligned}$$

and for  $0 < \alpha < 1$

$$\begin{aligned} |G_{\eta_0}^-| &\leq \frac{C_4(\alpha)}{(\eta_0 - \eta)^\beta} \left\{ \frac{|Y|}{\eta_0 - \eta} + \frac{1}{(2 - \xi_0 - \eta_0)^2} \right. \\ &\quad \left. + \frac{|Y|(\eta - \xi)}{(\xi_0 - \xi)(\eta_0 - \eta)} \left( \frac{\xi_0 - \eta}{\eta_0 - \eta} \right)^{1-\alpha} \right\} \sum_{i=1}^n |Y|^{i-1} \\ &\leq \frac{3C_4(\alpha)}{(\eta_0 - \eta)^\beta} \sum_{i=1}^n (2 - \xi_0 - \eta_0)^{-i-1}. \end{aligned}$$

Therefore, taking  $\alpha = \beta \in (0, 1)$ , we obtain estimates (A.29) and (A.31). □

### Appendix B: Auxiliary results

**Lemma B.1** *Suppose  $0 < \beta < 1$  and  $0 < \xi_0 < \eta_0 < 1$ . Then*

$$I^1(\xi_0, \eta_0) := \int_0^{\xi_0} \int_{\xi}^{\eta_0} H(\xi, \eta; \xi_0, \eta_0) d\eta d\xi \leq k_1 \xi_0. \tag{B.1}$$

*Proof* From (A.19) and (A.20) we obtain

$$\begin{aligned} I^1 &\leq C_H \left\{ \int_0^{\xi_0} \int_{\xi}^{\xi_0} (\xi_0 - \eta)^{-\beta} d\eta d\xi + \int_0^{\xi_0} \int_{\xi_0}^{\eta_0} (\eta - \xi_0)^{-\beta} d\eta d\xi \right\} \\ &=: C_H \{I_1^1 + I_2^1\}. \end{aligned}$$

Now we obtain

$$I_1^1 = \frac{\xi_0^{2-\beta}}{(1-\beta)(2-\beta)} \tag{B.2}$$

and

$$I_2^1 = \frac{1}{1-\beta} \xi_0(\eta_0 - \xi_0)^{1-\beta}. \tag{B.3}$$

Therefore estimate (B.1) holds. □

**Lemma B.2** *Suppose  $0 < \beta < 1$  and  $0 < \xi_0 < \eta_0 < 1$ . Then*

$$I^2(\xi_0, \eta_0) := \int_0^{\xi_0} \int_{\xi}^{\eta_0} |H_{\eta_0}(\xi, \eta; \xi_0, \eta_0)| d\eta d\xi \leq k_2 \xi_0 (\eta_0 - \xi_0)^{-\beta}. \tag{B.4}$$

*Proof* From (A.21) and (A.22) we obtain

$$\begin{aligned}
 I^2 &\leq C_H \left\{ \int_0^{\xi_0} \int_{\xi}^{\xi_0} \frac{(\xi_0 - \eta)^{-\beta}}{\eta_0 - \eta} d\eta d\xi + \int_0^{\xi_0} \int_{\xi_0}^{\eta_0} \frac{(\eta - \xi_0)^{-\beta}}{\eta_0 - \xi_0} d\eta d\xi \right\} \\
 &=: C_H \left\{ I_1^2 + \frac{I_2^1}{\eta_0 - \xi_0} \right\}.
 \end{aligned}$$

In  $I_1^2$  we substitute  $\eta = \xi + (\xi_0 - \xi)\sigma$  and, according to (A.2), we get

$$\begin{aligned}
 I_1^2 &= \int_0^{\xi_0} \left[ \int_0^1 (1 - \sigma)^{-\beta} \left( 1 - \frac{\xi_0 - \xi}{\eta_0 - \xi} \sigma \right)^{-1} d\sigma \right] \frac{(\xi_0 - \xi)^{1-\beta}}{\eta_0 - \xi} d\xi \\
 &= \frac{\Gamma(1)\Gamma(1 - \beta)}{\Gamma(2 - \beta)} \int_0^{\xi_0} F(1, 1, 2 - \beta; \zeta) \frac{(\xi_0 - \xi)^{1-\beta}}{\eta_0 - \xi} d\xi,
 \end{aligned}$$

where  $\zeta = \frac{\xi_0 - \xi}{\eta_0 - \xi}$ . Since  $c - a - b = -\beta < 0$ , according to (A.5), the hypergeometric function  $|F| \leq \text{const} \cdot (1 - \zeta)^{-\beta}$ . Therefore

$$I_1^2 \leq c_1(\eta_0 - \xi_0)^{-\beta} \int_0^{\xi_0} \left( \frac{\xi_0 - \xi}{\eta_0 - \xi} \right)^{1-\beta} d\xi \leq c_1 \xi_0 (\eta_0 - \xi_0)^{-\beta}. \tag{B.5}$$

Now (B.3) and (B.5) give estimate (B.4). □

**Lemma B.3** *Suppose  $0 < \beta < 1$  and  $0 < \xi_0 < \eta_0 < 1$ . Then*

$$I^3(\xi_0, \eta_0) := \int_0^{\eta_0} H(0, \eta; \xi_0, \eta_0) d\eta \leq k_3 \eta_0. \tag{B.6}$$

*Proof* Using (A.23) with  $\alpha = \beta$ , we obtain

$$\begin{aligned}
 I^3 &\leq c(\beta)\eta_0^{-\beta} \left\{ \int_0^{\xi_0} \frac{\eta^{2\beta}}{(\xi_0 - \eta)^\beta} d\eta + \int_{\xi_0}^{\eta_0} \frac{\eta^{2\beta}}{(\eta - \xi_0)^\beta} d\eta \right\} \\
 &\leq c(\beta)\eta_0^{-\beta} \left\{ \xi_0^{2\beta} \int_0^{\xi_0} (\xi_0 - \eta)^{-\beta} d\eta + \eta_0^{2\beta} \int_{\xi_0}^{\eta_0} (\eta - \xi_0)^{-\beta} d\eta \right\} \\
 &= \frac{c(\beta)}{1 - \beta} \eta_0^{-\beta} \left\{ \xi_0^{1+\beta} + \eta_0^{2\beta} (\eta_0 - \xi_0)^{1-\beta} \right\} \\
 &\leq k_3 \eta_0. \tag{B.6}
 \end{aligned}$$

□

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the writing of this paper. The authors read and approved the final manuscript.

**Author details**

<sup>1</sup>Faculty of Mathematics and Informatics, University of Sofia, Sofia, 1164, Bulgaria. <sup>2</sup>Faculty of Applied Mathematics and Informatics, Technical University of Sofia, Sofia, 1000, Bulgaria. <sup>3</sup>Faculty of Mathematics, Karlsruhe Institute of Technology, Karlsruhe, 76131, Germany.

**Acknowledgements**

The authors thank the anonymous referees for making several helpful suggestions. The research of Tsvetan Hristov, Aleksey Nikolov and Nedyu Popivanov was partially supported by the Sofia University Grant 152/2016.

## References

- Morawetz, C: Mixed equations and transonic flow. *J. Hyperbolic Differ. Equ.* **1**(1), 1-26 (2004)
- Protter, M: New boundary value problem for the wave equation and equations of mixed type. *J. Ration. Mech. Anal.* **3**, 435-446 (1954)
- Morawetz, C: A weak solution for a system of equations of elliptic-hyperbolic type. *Commun. Pure Appl. Math.* **11**, 315-331 (1958)
- Lax, P, Phillips, R: Local boundary conditions for dissipative symmetric linear differential operators. *Commun. Pure Appl. Math.* **13**, 427-455 (1960)
- Aziz, A, Schneider, M: Frankl-Morawetz problems in  $R^3$ . *SIAM J. Math. Anal.* **10**, 913-921 (1979)
- Barros-Neto, J, Gelfand, I: Fundamental solutions for the Tricomi operator. *Duke Math. J.* **128**(1), 119-140 (2005)
- Dechevski, L, Popivanov, N: Morawetz-Protter 3D problem for quasilinear equations of elliptic-hyperbolic type. Critical and supercritical cases. *C. R. Acad. Bulgare Sci.* **61**(12), 1501-1508 (2008)
- Keyfitz, B, Tesdall, A, Payne, K, Popivanov, N: The sonic line as a free boundary. *Q. Appl. Math.* **71**(1), 119-133 (2013)
- Lupo, D, Monticelli, D, Payne, K: On the Dirichlet problem of mixed type for lower hybrid waves in axisymmetric cold plasmas. *Arch. Ration. Mech. Anal.* **217**(1), 37-69 (2015)
- Lupo, D, Morawetz, C, Payne, K: On closed boundary value problems for equations of mixed elliptic-hyperbolic type. *Commun. Pure Appl. Math.* **60**(9), 1319-1348 (2007)
- Lupo, D, Payne, K, Popivanov, N: Nonexistence of nontrivial solutions for supercritical equations of mixed elliptic-hyperbolic type. *Prog. Nonlinear Differ. Equ. Appl.* **66**, 371-390 (2006)
- Lupo, D, Payne, K, Popivanov, N: On the degenerate hyperbolic Goursat problem for linear and nonlinear equations of Tricomi type. *Nonlinear Anal.* **108**, 29-56 (2014)
- Rassias, J: Tricomi-Protter problem of nD mixed type equations. *Int. J. Appl. Math. Stat.* **8**(M07), 76-86 (2007)
- Kalmenov, T, Koshanov, B, Iskakova, U: Completeness of the root vectors to the Tricomi problem for Lavrent'ev-Bitsadze equation. In: *Nonclassical Equations of Mathematical Physics. Proceedings of the Seminar Dedicated to the 60-th Birthday of Prof. V. N. Vragov, Novosibirsk, Russia*, pp. 125-129. Sobolev Institute of Mathematics RAN, Novosibirsk (2005)
- Moiseev, T: Gellerstedt problem with a generalized Frankl matching condition on the type change line with data on external characteristics. *Differ. Equ.* **52**, 240-247 (2016)
- Sadybekov, M, Yessirkegenov, N: Properties of solutions to Neumann-Tricomi problems for Lavrent'ev-Bitsadze equations at corner points. *Electron. J. Differ. Equ.* **2014**, 193 (2014)
- Otway, T: *The Dirichlet Problem for Elliptic-Hyperbolic Equations of Keldysh Type. Lecture Notes in Mathematics*, vol. 2043. Springer, Berlin (2012)
- Otway, T: *Elliptic-Hyperbolic Partial Differential Equations. A Mini-Course in Geometric and Quasilinear Methods. Springer Briefs in Mathematics*. Springer, Cham (2015)
- Otway, T, Marini, A: Strong solutions to a class of boundary value problems on a mixed Riemannian-Lorentzian metric. *Discrete Contin. Dyn. Syst.* **2015**(Suppl.), 801-808 (2015)
- Oleinik, O, Radkevič, E: *Second Order Equations with Nonnegative Characteristic Form*. Am. Math. Soc., Providence (1973)
- Nakhushiev, A: *Problems with Shifts for Partial Differential Equations*. Nauka, Moscow (2006) [in Russian]
- Chen, S: A mixed equation of Tricomi-Keldysh type. *J. Hyperbolic Differ. Equ.* **9**(3), 545-553 (2012)
- Čanić, S, Keyfitz, B: A smooth solution for a Keldysh type equation. *Commun. Partial Differ. Equ.* **21**(1-2), 319-340 (1996)
- Marin, M, Lupu, M: On harmonic vibrations in thermoelasticity of micropolar bodies. *J. Vib. Control* **4**(5), 507-518 (1998)
- Marin, M, Craciun, E-M, Pop, N: Considerations on mixed initial-boundary value problems for micropolar porous bodies. *Dyn. Syst. Appl.* **25**(1-2), 175-196 (2016)
- Protter, M: A boundary value problem for the wave equation and mean value problems. *Ann. Math. Stud.* **33**, 247-257 (1954)
- Kalmenov, T: On the strong solutions of the Darboux and Tricomi problems. *Sov. Math. Dokl.* **28**, 672-674 (1983)
- Nakhushiev, A: Criteria for continuity of the gradient of the solution to the Darboux problem for the Gellerstedt equation. *Differ. Equ.* **28**(10), 1445-1457 (1992)
- Garabedian, P: Partial differential equations with more than two variables in the complex domain. *J. Math. Mech.* **9**, 241-271 (1960)
- Khe, KC: On nontrivial solutions of some homogeneous boundary value problems for the multidimensional hyperbolic Euler-Poisson-Darboux equation in an unbounded domain. *Differ. Equ.* **34**(1), 139-142 (1998)
- Popivanov, N, Schneider, M: The Darboux problems in  $R^3$  for a class of degenerating hyperbolic equations. *J. Math. Anal. Appl.* **175**(2), 537-579 (1993)
- Bitsadze, A: *Some Classes of Partial Differential Equations*. Gordon & Breach, New York (1988)
- Hristov, T, Popivanov, N: Singular solutions to Protter's problem for a class of 3-D weakly hyperbolic equations. *C. R. Acad. Bulgare Sci.* **60**(7), 719-724 (2007)
- Popivanov, N, Popov, T: Asymptotic expansions of singular solutions for  $(3 + 1)$ -D Protter problems. *J. Math. Anal. Appl.* **331**, 1093-1112 (2007)
- Nikolov, A, Popivanov, N: Asymptotic expansion of singular solutions to Protter problem for  $(2 + 1)$ -D degenerate wave equation. In: *Venkov, G, Pasheva, V (eds.) Applications of Mathematics in Engineering and Economics: AMEE'13, Sozopol, Bulgaria, June 2013. AIP Conf. Proc.*, vol. 1570, pp. 249-256. AIP, New York (2013)
- Popivanov, N, Popov, T: Behaviour of singular solutions to 3-D Protter problem for a degenerate hyperbolic equation. *C. R. Acad. Bulgare Sci.* **63**(6), 829-834 (2010)
- Nikolov, A, Popivanov, N: Singular solutions to Protter's problem for  $(3 + 1)$ -D degenerate wave equation. In: *Venkov, G, Pasheva, V (eds.) Applications of Mathematics in Engineering and Economics: AMEE'12, Sozopol, Bulgaria, June 2012. AIP Conf. Proc.*, vol. 1497, pp. 233-238. AIP, New York (2012)
- Didenko, V: On boundary-value problems for multidimensional hyperbolic equations with degeneration. *Sov. Math. Dokl.* **13**, 998-1002 (1972)

39. Aldashev, S: A criterion for the existence of eigenfunctions of the Darboux-Protter spectral problem for degenerating multidimensional hyperbolic equations. *Differ. Equ.* **41**, 833-839 (2005)
40. Hristov, T, Popivanov, N, Schneider, M: On the uniqueness of generalized and quasi-regular solutions for equations of mixed type in  $R^3$ . *Sib. Adv. Math.* **21**(4), 262-273 (2011)
41. Hristov, T, Popivanov, N, Schneider, M: Generalized solutions to Protter problems for 3-D Keldysh type equations. In: Sivasundaram, S (ed.) 10th International Conference on Mathematical Problems in Engineering, Aerospace and Sciences: ICNPAA 2014, Narvik, Norway, July 2014. AIP Conf. Proc., vol. 1637, pp. 422-430. AIP, New York (2014)
42. Hristov, T: Singular solutions to Protter problem for Keldysh type equations. In: Venkov, G, Pasheva, V (eds.) Applications of Mathematics in Engineering and Economics: AMEE'14, Sozopol, Bulgaria, June 2014. AIP Conf. Proc., vol. 1631, pp. 255-262. AIP, New York (2014)
43. Hristov, T, Popivanov, N, Schneider, M: Protter problem for 3-D Keldysh type equations involving lower order terms. In: Applications of Mathematics in Engineering and Economics: AMEE'15, Sozopol, Bulgaria, June 2015. AIP Conf. Proc., vol. 1690, pp. 04002001-04002012. AIP, New York (2015)
44. Hristov, T, Nikolov, A, Popivanov, N, Schneider, M: Generalized solutions of Protter problem for  $(3 + 1)$ -D Keldysh type equations. In: Pasheva, V, Venkov, G, Popivanov, N (eds.) Applications of Mathematics in Engineering and Economics: AMEE'16, Sozopol, Bulgaria, June 2016. AIP Conf. Proc., vol. 1789, pp. 04000701-04000713. AIP, New York (2016)
45. Jones, M: Spherical Harmonics and Tensors for Classical Field Theory. Research Studies Press, Letchworth (1986)
46. Gellerstedt, S: Sur une équation linéaire aux dérivées partielles de type mixte. *Ark. Mat. Astron. Fys. A* **25**(29), 1-23 (1937)
47. Smirnov, M: Degenerating Hyperbolic Equations. Vysheishaia shkola Publ. House, Minsk (1977) [in Russian]
48. Moiseev, E: Approximation of the classical solution of a Darboux problem by smooth solutions. *Differ. Equ.* **20**, 59-74 (1984)
49. Bateman, H, Erdelyi, A: Higher Transcendental Functions, vol. 1. McGraw-Hill, New York (1953)
50. Smirnov, M: Mixed Type Equations. Vishaya shkola Publ. House, Moskow (1985) [in Russian]
51. Volkodavov, V, Zaharov, V: Tables of Riemann and Riemann-Hadamard Functions for Some Differential Equations in  $n$ -Dimensional Euclidean Spaces. Samara State Teacher's Training University, Samara (1994) [in Russian]
52. Meredov, M: The unique solvability of Darboux's problem for a degenerating system. *Differ. Equ.* **10**, 63-70 (1975)

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---